



NASA Conference Publication 2231

Controlled Ecological Life Support System

Use of Higher Plants

NASA-CP-2231 19820016958

*Proceedings of two NASA workshops held at
the O'Hare Airport Conference Center
Chicago, Illinois, November 1979
and at the Ames Research Center
Moffett Field, California
March 1980*

NASA

NASA Conference Publication 2231

Controlled Ecological Life Support System

Use of Higher Plants

Edited by
T. W. Tibbits
University of Wisconsin
Madison, Wisconsin

D. K. Alford
Metropolitan State College
Denver, Colorado

Proceedings of two NASA workshops held at
the O'Hare Airport Conference Center
Chicago, Illinois, November 1979
and at the Ames Research Center
Moffett Field, California
March 1980

NASA

National Aeronautics
and Space Administration

**Scientific and Technical
Information Branch**

1982

CONTENTS

	<u>Page</u>
PREFACE	v
I. INTRODUCTION	1
II. CRITERIA FOR PLANT SELECTION	
A. Food Production	13
B. Nutrition	13
C. Oxygen Production and Carbon Dioxide Utilization	18
D. Water Recycling	21
E. Waste Recycling	21
F. Other Criteria	22
III. LITERATURE CITED	23
IV. RECOMMENDATIONS	
A. Plant Species	27
B. Growing Procedures	31
C. Research Priorities	35
V. BIBLIOGRAPHY AND PRODUCTIVITY OF THE SELECTED PLANT SPECIES	39
VI. WORKSHOP PARTICIPANTS	80

PREFACE

This report has been developed by a group of 27 crop physiologists that met in two separate workshops held in Chicago, IL, November 1979 and Moffett Field, CA, March 1980. The recommendations and conclusions presented in this report represent a unified consensus of this group.

The purpose of the workshops was to consider the use of higher plants in Controlled Ecological Life Support Systems (CELSS). The results of the workshops are intended to contribute to the development of a comprehensive program plan for NASA's Biological Systems Research program.

The excellent discussions that ensued and the harmonious accord that was obtained at the workshops were due in large measure to the following physiologists who provided direction and chaired separate sessions: Johan Hoff, Tak Hoshizaki, Bob Langhans, Doug Ormrod, Ralph Prince, Dave Raper, Frank Salisbury and Herb Ward. A special thanks to Ben Zietman and Pete Zill for logistics and coordination at NASA/Ames. These workshops were supported by NASA under Grant No. NSG 2405.

I. INTRODUCTION

The use of higher plants for food production has been proposed for long duration or large scale manned space missions to minimize the prohibitively large storage and resupply costs (Mason and Carden 1979; Spurlock et al. 1979; Johnson and Holbrow 1977) associated with carry-on food and oxygen. There is no consensus on the break-even point (weight) in flight duration for a contained life support system compared with food storage or resupply alternatives. The break even point has been variously described as anywhere from 30 days to 12 years (Space Science Board 1969; Ward et al. 1963). Estimates vary widely due to incomplete analyses and well recognized difficulties encountered in extrapolation from theoretical models and experimental data (Ward et al. 1963). A major deficiency has been lack of data for maximum food producing capability of plants and the minimum weight requirements for maintaining plants in sustained growth systems.

Initial efforts toward development of food production systems for manned spacecraft were initiated by both the United States and the Soviet Union during the late 1950's and early 1960's. Systems were planned for both orbital and moon-based stations (Taub, 1974). Significant funds were committed to study the use of different kinds of life support systems, including algae, particularly Chlorella (Dole 1964; Drake 1966; Gitelson et al. 1975; Miller and Ward 1966; Ward et al. 1963), Hydrogenomonas (Drake 1966), various species of duckweed (Ward et al. 1963), and physiochemical systems lacking biological components (Dole 1964). Major emphasis was placed on developing continuous production algal systems, since they use growing space efficiently, produce oxygen, are rich in protein, may be used as food supplements, and are also efficient in processing metabolic wastes. However, productive algal systems have been difficult to maintain for long periods of time, do not provide a balanced palatable diet and require cumbersome maintenance and harvesting procedures (Dole 1964).

It has been proposed by the Space Science Board of the National Academy of Sciences (1969) that multiorganism systems (plants and algae) should be employed for food and oxygen production in future space efforts. This would provide two support systems, so that if one failed or was inoperative for a period of time, production of both food and oxygen would continue.

Higher Plants

Higher plants can provide most, if not all, of the major food needs of man; calories, proteins, fats and carbohydrates; along with the much smaller requirements for minerals, vitamins and trace elements (Table 1). The nutritional value of the major U.S. plant foods expressed on a 100 g edible portion is indicated in Tables 2a, 2b and 2c. The nutritional value contained in individual servings in Tables 3a and 3b. In addition, higher plants can provide all of the oxygen required for life support in manned spacecraft and recycle waste water. The daily food, oxygen and water requirements for man are shown in Table 4. It has been variously estimated that anywhere from 4m^2 (Dadykin 1968) to 250m^2 (Dole 1964) of hydroponically grown plants are required to provide the life support requirements for one man.

Heydecker (1973) expressed concern that plants produce seeds with low viability when grown for several generations in controlled environments and that the edible portions will not be as nutritious or as productive as field grown plants. However, Gitel'son et al. (1975) reported that cultivars reproduced up to 10 generations of healthy plants in controlled environments without deterioration of vigor. Similarly, Gitel'son et al. (1975) compared protein, fat, vitamin and mineral content of field and phytotron grown beets, radishes, turnips and onions and found in all cases, nutritional levels in phytotron-grown plants were comparable or better than field-grown plants (Table 5). It is generally recognized by crop physiologists that the nutritive value of plants grown in controlled environments varies considerably and has nutritive value that is simi-

Table 1

DAILY DIETARY RECOMMENDATIONS AND PRACTICES

NUTRIENT	NRC ¹ (RDA)	FAO/WHO ²	SKYLAB ³	Veg. ⁴
Energy (kcal)	2700 (2300-3100)	3000	2700 (2300-3100)	1970
Protein (g)	56	37-62	90-25±10	65.4
Vit. A (mcg R.E.)	1000	750	1000	2102
Vit. D (mcg)	5 (200 I.U.)	100 I.U.	5 (200 I.U.)	-
Vit. E (mg Alpha T.E.)	10	-	10	-
Vit. C (mg)	60	30	60	180
Thiamin (mg)	1.4	1.2	1.4	1.9
Riboflavin (mg)	1.6	1.7	1.6	1.2
Niacin (mg)	18	19.8	18	18
Vit. B ₆ (mg)	2.2	-	2.2	-
Folacin (mg)	400	200	400	-
Vit. B ₁₂ (mcg)	3	2	3	0
Calcium (mg)	800	400-500	750-850±16	594
Phosphorus (mg)	800	-	1500-1700±120	1368
Magnesium (mg)	350	-	300-400±100	-
Iron (mg)	10 (man) 18 (woman)	9 (man) 28 (woman)	10 (man) -	19
Zinc (mg)	15	-	15	-
Iodine (mcg)	150	-	150	-
Sodium (g)	1.1-3.3		3.0-6.0±0.5	2.2
Potassium (mg)	1525-4575		2740 min. no max. and no range	4100

¹Reference man 70 kg mixed diet (Nat. Acad. Sci. 1980)

²Reference man 65 kg mixed diet (FAO 1967, FAO 1970b, FAO 1973)

³Mixed diet (Klicka *et al.* 1967)

⁴VEG: Average calculated values of a 14-day vegetarian cycle menu which was developed from communication with practicing strict vegetarians; and adapted from "Recipes from a Small Planet" (Robertson *et al.* 1978) and "Laurel's Kitchen, a handbook for Vegetarian Cookery and Nutrition" (Ewald 1977). Soybean meal values were substituted for milk and milk products. Soybean lecithin values were used in place of egg in mayonaise, etc.

Table 2a

COMPOSITION OF SELECTED PLANT SPECIES¹

General

(Amount in 100g Edible Portion-Dried)

<u>Species</u>	<u>Cal- ories</u>	<u>Prot</u>	<u>Fat</u>	<u>Ash</u> (grams)	<u>CHO</u>	<u>Fiber</u>
Soybean	466	37	19.5	5.0	38	8.6
Dry Bean	386	24	1.8	4.4	70	4.5
Split Pea	382	27	1.1	3.1	69	1.3
Podded Pea	317	20	1.2	6.6	72	7.2
Chick Pea	403	23	5.4	3.4	68	5.6
Peanut	585	28	45	2.8	24	2.5
Rice	409	8.5	1.9	1.3	88	.7
Wheat	379	16	2.5	2.0	79	2.6
Oats	425	16	8.1	2.1	74	1.3
Corn	411	10	3.9	1.3	85	1.1
Potato	373	9	.5	5.4	86	1.8
Sweet Potato	290	6	2.2	3.5	89	3.2
Beet Greens	338	13	.8	8.9	77	7.3
Lettuce	288	23	3.8	17.3	56	11.5
Spinach	274	32	4.1	20.5	44	8.2
Mustard Greens	295	29	4.8	13.3	53	10.5
Kale	298	29	4.5	12.7	54	9.0
Tomato	339	17	5.1	10.2	68	10.2
Strawberry	366	7.9	5.0	5.0	82	13.9
Onion	366	11.4	1.6	4.9	84	0.8
Cucumber	308	17.9	2.6	10.3	69	12.8
Broccoli	287	32.7	2.0	10.9	54	12.9
Sugar Beet	320				80	4.0

¹Proximate analysis obtained from Watt and Merrill (1963).

Table 2b

COMPOSITION OF SELECTED PLANT SPECIES¹

Vitamins

(Amount in 100g Edible Portion-Raw)

	A (IU)	B ₁ (mg)	B ₂ (mg)	B ₇ (mg)	B ₆ (mg)	Fol (µg)	Pan (mg)	Bio (µg)	C (mg)	E (mg)	Car (mg)
Soybean	80	1.1	.3	2.2							
Dry Bean		.7	.2	2.4	.6		.7			2.3	
Split Pea		.7	.2	3.2	.1	33	2.0				
Podded Pea		.2	.1	.9	.2	25	.8		36	2.7	
Chick Pea		.5	.15	1.5		180					190
Peanut		.9	.1	16	10.5	110	2.7		3	8.1	
Rice		.3	.05	4.7							
Wheat		.5	.1	4.7	.6		.8			2.3	
Oats		.5	.1	1.0	.1	60	1.0	20		.8	
Corn	510	.4	.1	2.0	(.2) ²	(52)	(.5)			(.8)	(240)
Potato		.1	.04	1.2	.2	10	.2		30	.1	
Sweet Potato	8800	.1	.06	.6					21		
Beet Greens	6100	.1	.2	.4					30		
Lettuce	1900	.05	.08	.4					18		
Spinach	8100	.1	.2	.6					51		
Mustard Greens	7000	.1	.2	.8					97		
Kale	8900	(.2)	(.3)	(2.0)					(125)		
Tomato	900	.1	.04	.7					23		
Strawberry	60	.03	.07	.6					59		
Onion	40	.03	.04	.2					10		
Cucumber	250	.03	.04	.2					11		
Broccoli	2500	.1	.2	.9					113		
Sugar Beet											

¹Obtained from Watt and Merrill (1963).²Estimated values.

Table 2c

COMPOSITION OF SELECTED PLANT SPECIES¹

Minerals

(Amount in 100g Edible Portion-Raw)

Species	Ca	Mg	Na	K	Cu	Zn	Fe	S	Cl	P
	(milligrams)									
Soybean	226	265	5	1677			8.4			554
Dry Bean	144	170	19	1196	.85	.20	7.8			425
Pea, Split	33	180	38	910	.58	(4.0) ²	5.4	170	56	270
Pea and Pod	49	35	9	135	.22	(1.3)	1.1			54
Chick Pea	140	160	40	800	.76		6.4	180	60	300
Peanut	61	180	6	680	.27	3.0	2.0	380	7	370
Rice, Brown	32	88	9	214			1.6			221
Wheat, Whole	36	.9	(3)	435	.20		3.1		.4	383
Oats	55	110	33	370	.23	(3.0)	4.1	160	73	380
Corn	22	147	1	284			2.1			268
Potato, White	8	34	7	570	.15	.3	.5	35	79	40
Potato, Sweet	32	31	10	243			.7			47
Beet, Greens	119	106	(130)	570			3.3			40
Lettuce	68	(11)	9	264	(.03)	.2	1.4		53	25
Spinach	93	88	71	470			3.1			51
Mustard Greens	183	27	32	377	.12		3.0	170	89	50
Kale	179	37	75	378			2.2			73
Tomato	13	14	3	244			.5			27
Strawberry	21	12	1	164			1.0			21
Onion	27	12	10	157			.5			36
Cucumber	25	11	6	160			1.1			27
Broccoli	103	24	15	382			1.1			78
Sugar Beets										

¹Obtained from Watt and Merrill (1963) and Paul and Southgate (1978).²Estimated values.

Table 3a

COMPOSITION OF SELECTED PLANT SPECIES¹

General

Species No.	Cal- ories	(per serving)					Serving Size	Cooking Method
		Prot (g)	Fat (g)	Ash (g)	CHO (g)	Fiber (g)		
Soybean	117	9.9	5.2	1.4	9.7	2.3	1/2 cup = 90g	boiled
Dry Bean	106	7.0	.5	1.3	19.1	1.3	1/2 cup = 95g	boiled
Split Pea	118	9.3	.4	1.1	24	.5	1/2 cup = 100g	boiled
Podded Pea	(50) ²	(3)	(.2)	(1)	(13)	(1.3)	(1/2 cup)	boiled
Chickpea	(113)	(9)	(2)	(1)	(24)	(2)	(1/2 cup)	boiled
Peanut	154	7.3	11.9	.7	6.4	.6	27g	roasted
Rice	116	2.5	.6	.4	25	.2	1/2 cup = 98g	boiled
Wheat	114	4.2	.5	(.6)	24	.8	1/2 cup = 68g	parboiled bulgur
Oats	66	2.4	1.2	.3	12	.2	1/2 cup = 120g	boiled oatmeal
Corn	68	1.7	.6	.2	14	.2	1/3 cup = 55g	whole grain boiled
Potato	84	2.0	.1	1.2	19	.4	1 tuber = 101g	boiled
Sweet Potato	252	3.7	1.4	2.3	57	2.1	1 = 205g	boiled
Beet Greens	13	1.3	.2	(1.0)	2.4	0.7	1/2 cup = 73g	boiled
Lettuce	8	.6	.1	.5	1.5	.3	2 lg. or 4 sm. leaves = 50g	raw
Spinach	21	2.5	.5	1.5	2.9	.8	1 cup = 80g	boiled
Mustard Greens	15	1.6	.2	.8	2.8	.6	1/2 cup = 70g	boiled
Kale	15	1.6	.2	.8	2.8	.6	1/2 cup = 70g	boiled
Tomato	30	1.5	.5	.9	6.0	.9	1 med = 150g	raw
Strawberry	55	1.2	.8	.8	12.1	2.1	1 cup = 149g	raw
Onion	19	.5	.1	.3	4.4	.4	50g	boiled
Cucumber	6	.4	.1	.2	1.4	.3	50g	raw
Broccoli	15	1.7	.1	.6	2.8	.7	1 cup = 150g	boiled
Sugar Beet								

¹Proximate analysis obtained from Adams (1975).²Estimated values.

Table 3b

COMPOSITION OF SELECTED PLANT SPECIES¹

Vitamins

(per serving)

	A IU	B ₁ mg	B ₂ mg	Nia mg	B ₆ mg	Fd. μg	Pan. mg	Bio μg	C mg	E mg	Car. μg	Serving
Soybean	25	.2	.08	0.6							(50)	1/2 cup = 90g
Dry Bean	0	.14	.07	.7								1/2 cup = 85g
Split Pea	40	.15	.04	.9								1/2 cup = 100g
Podded Pea	(500) ²	(.2)	(.07)	(1.6)								1/2 cup
Chickpea	50	.31	.15	2.0		37					210	1/2 cup = 100g
Peanut		.23	.19	12.4	.3		1.4			22		1/2 cup = 72g
Rice	0	.09	.04	2.7	(.05)	(6)	(.2)	(1.0)		(.1)		1/2 cup = 83g
Wheat	0	.33	.07	2.6								1/2 cup = 60g
Oats	0	.10	.03	.1								1/2 cup = 120g
Corn	200	.12	.04	.8	.09	18	.21			.3	13	1/2 cup = 58g
Potato	tr.	.12	.05	2.0	.18	10	.20		22	.1		1 = 150g
Sweet Potato	11940	.14	.09	.9					26			1 = 180g
Beet Greens	3700	.05	.11	.2					11			1/2 cup = 73g
Lettuce	1050	.03	.04	.2					10			55g
Spinach	7300	.07	.13	.5					25			1/2 cup = 90g
Mustard Greens	4060	.06	.10	.4					34			1/2 cup = 70g
Kale	4600	.06	.10	.9					51			1/2 cup = 55g
Tomato	410	.05	.04	.6					21			1 = 100g
Strawberry	90	.04	.10	.9					88			1 cup = 149g
Onion	40	.03	.03	.2					8			1/2 cup = 105g
Cucumber	130	.02	.02	.1					6			1/2 cup = 53g
Broccoli	3500	.13	.28	1.1					126			1 stalk = 140g
Sugar Beet												

¹Obtained from Adams (1975).²Estimated value.

Table 4

DAILY LIFE SUPPORT REQUIREMENTS

Type of Input	Amounts (kg/person/day)		
	Gitel'son 1975	Modell 1977	Ward and Miller 1966
Food (dry)	0.6		0.52
Oxygen	0.9		0.86
Drinking water	1.8	2.2-2.5	2.2
Sanitary water	2.3	6.5	
Domestic water	16.8		

Table 5

A COMPARISON OF NUTRITION LEVELS IN FIELD AND PHYTOTRON
GROWN RADISHES¹

	<u>Field</u>	<u>Phytotron</u>
total amino acid content (%)	8.67	9.50
essential amino acid content (%)	3.56	3.61
mineral content (% dry matter)	7.89	8.25
ascorbic acid (mgm %)	21.4	37.6

¹Obtained from Gitel'son 1975.

lar to that of plants grown in field environments.

Thus far, there have been few growth studies of higher plants conducted within spacecraft. The United States biosatellite experiments studied the effects of zero gravity on wheat seedlings and pepper plants over several days (Johnson and Tibbitts 1968; Lyon 1968). Russian studies in space have concentrated upon study of effects of radiation on plants. Some chromosomal aberrations have been found (Glembotskiy et al. 1962; Nikolayev et al. 1964). It should be noted, however, that the Apollo-Soyuz spacecraft carried Arabidopsis, Nicotiana and Zea but no significant differences were found between space grown plants and ground based plants (Anderson et al. 1979). Results indicated that plants tolerate radiation levels through processes of photo-adaption and repair injuries by photoactivation (Shakov et al. 1962) or may be protected by prior chemical treatment (Shaykarov 1965). The United States Space Science Board (1969) has concluded that there should be no significant radiation shielding problems in future space ventures. Therefore, radiation effects will not be considered further in this report.

There also is concern about the possible accumulation of toxins within a regenerative life support system, which might damage or kill plants or perhaps harm humans. Potential contaminants may be carried in the air, root medium or water. They may originate from man, the spacecraft, or the plants themselves. Plants are known to give off at least 200 discrete substances including hydrocarbons, aldehydes, alcohols, ketones, ethers (Gitel'son et al. 1975), carbon monoxide (Wilks 1959), various organic acids, amino acids, lactones, flavones (Dadykin 1968) as well as ethylene and terpenes that may be injurious and/or prevent normal plant growth (Ormrod 1979). Humans also release 150 volatile substances that may concentrate in a closed system and disrupt plant growth (Gitel'son et al. 1975). Plants such as beets give off

substances which might be harmful to other plants and even to the plant producing the substances (Milov et al. 1975). The significance of plant toxins was demonstrated in the BIOS III project conducted in the Soviet Union (Gitel'son et al. 1975). BIOS III was a six month, multiorganism study involving man, higher plants and algae systems. It was shown that higher plants grew vigorously without the algae system, but the plants died shortly after the algae were introduced into the unit. Presumably, unknown toxins given off by the algae killed the plants. Some chemicals given off by plants may accumulate within a regenerative life support system and be toxic to man. For example, radishes, tomatoes, carrots, and potatoes give off propionaldehyde which is toxic to humans (Rusakova et al. 1975).

Care should be taken to insure that seed stock and culture systems are reasonably free of pathogenic organisms. There is no evidence that a totally disease-free environment would be necessary or reasonable for effective plant growth in a closed system. Plants growing vigorously in controlled environments rarely develop disease problems unless pathogens are introduced by some external source.

Precise control of environmental conditions will be necessary to obtain maximum crop yields within a regenerative life support system. Assuming the availability of nuclear or solar energy (Miller and Ward 1966) space systems should not be constrained by energy availability. Therefore, continuous high irradiance lighting may be available. A few studies have demonstrated that plant growth is most rapid with continuous light of moderately high intensity (greater than $300 \mu\text{E m}^{-2} \text{sec}^{-1}$) (Dolon 1973; Rao Rama Rao 1965). However, most plants have been found to have better growth with alternating light and dark periods that provide maximum yields with 16-18 hour light and 6-8 hour dark (Evans 1969; Gitel'son et al. 1975; Kristoffersen 1963).

Russian scientists have proposed that the light level for effective use of

plants in a regenerative life support system should be at least $50-60 \text{ Wm}^{-2}$ PPFD ($175-200 \mu\text{E sec}^{-1}\text{m}^{-2}$ PPFD or 10-12 klux) (Gitel'son et al. 1975). However, it should be recognized that the rate of dry matter accumulation by most plants will increase with increasing light levels to 150 Wm^{-2} and higher for certain species (Warrington et al. 1978).

The spacing of plants within a growing system is critical for maximum yield. It has been established that the optimum leaf area index (vertical density of leaves) for planting varies depending on light intensity and quality of light. For instance, it has been estimated that a leaf index of 7-8 would be required for maximum yield when the level of radiation is 150 Wm^{-2} PPFD (Dadykin 1968).

One of the significant problems of plant growth in a controlled environment system is the selection of an appropriate rooting medium that will afford optimum nutrition and yet be practical in terms of total amount of salt and water required. Previous experiments proposed for space systems have detailed a variety of techniques, including liquid culture (Krauss 1962), aeroponics (Dadykin 1968; Milov et al. 1975), artificial nutrient media such as vermiculite (Anonymous 1965; Lebedeva 1964) and perlite (Milov et al. 1975). Other methods under study include the nutrient film technique (Dadykin 1968) and subirrigation air cultivation which has been used to successfully cultivate grain crops within regenerative life support systems (Gitel'son et al. 1975). The use of aeroponics and subirrigation air cultivation techniques can minimize significantly the total weight of the plant-culture system. However, these systems require frequent and regular care to maintain desirable salt balance and solution pH. Growth in artificial media also can result in the uptake of undesirable chemicals into plants. Problems have been reported for many different media. For example, it has been reported that plants can take up high level of fluorides if grown in perlite (Ormrod 1979).

II. CRITERIA FOR PLANT SELECTION

A. Food Production

Species and cultivars should be selected that produce the maximum quantity of digestible biomass and the minimum quantity of non-digestible biomass based on an integration of the following unit factors:

- a. volume of space required per unit time
- b. labor requirements
- c. weight of the plant-growing system
- d. electrical energy utilized
- e. purchase and maintenance costs of plant-growing system

Integration of these factors for each plant species will require determining the magnitude of each of these unit factors. Therefore, a primary requirement for the plant research for life support systems will be in determining these unit factors and developing mechanisms for minimizing their impact.

B. Nutrition

Plants grown in the CELSS must be able to provide a nutritionally and psychologically satisfactory diet for the human inhabitants. The selected plant species must be evaluated in terms of a number of "use" criteria, including: energy concentration, nutritional composition, palatability, processing requirement, acceptable serving size and frequency, flexibility of usage, storage stability, toxicity, degree of human nutritional experience. A group of plant species should be selected that will provide balanced nutrition at levels sufficient to cover requirements for humans. However, to obtain maximum efficiency, the possibility of supplementing the diet with animal protein, and certain prepared minerals and vitamins should be included in diet development. The efficiency of different plant species should be compared on the basis of the quantity of food that is digested by humans.

It is possible for an adult person to obtain sufficient energy on a strict vegetarian ("vegan") diet. The amount of energy required is related to age and weight (Table 6), sex, degree of physical activity, including physical and psychological stress, and environmental conditions. For an average adult diet of 2300 calories, about 400 grams of carbohydrates per day should be included (Wilks 1964). The U.S. Dietary Goals suggests that the distribution of energy sources for the general public as 58% from carbohydrates, 12% from protein, and 30% from fat. A somewhat higher energy contribution from protein is recommended under stressfull conditions (Gemini and Appollo projects, Table 7). Sugar usage should be limited to adding culinary interest and/or to increase energy intake whenever this is difficult through other means.

Protein is of major concern in a vegetarian diet both for energy and to supply needed amino acids for the body. Plant proteins occur generally at low concentration, except in legume seed and nuts and their quality is generally poor. Protein, from different plants, vary in the type and proportion of amino acids. Ideally, amino acids should be absorbed from the digestive tract of humans in proportions similar to their occurence in the tissues that are being replaced. The amino acid composition of individual plant proteins are compared to that of the human body or whole egg protein, which often is taken as a reference protein (Table 8). It is generally found that most plant proteins are not fully utilized in the intestinal tract. The digestibility coefficient varies from 20% for sorghum to 89% for certain wheat flours. Most plant proteins have a digestibility coefficient fairly close to 75%. When a meal is composed of proteins from several sources and has a total composition more closely resembling the requirements of essential amino acids (Table 9), it is found that the digestibility is also improved. The diet should contain between 50-100 grams of protein (Table 1).

Table 6
CALORIC REQUIREMENT OF ADULT MALES
PER DAY¹ (Normal activity)

Body Wt.		Kilocalories at Age:		
		25	45	65
kg.	(lb)			
50	110	2300	2050	1750
60	132	2600	2350	1950
70	154	2900	2600	2200
80	176	3200	2900	2450

¹Adapted from Consolazio (1964).

Table 7
CALORIE INTAKE PATTERNS OF DAILY DIETS¹

Food Constituent	Space Programs				
	U.S. Current Practice	U.S. Dietary Goal	Project Gemini (2500 kcal)	Project Apollo (2800 kcal)	Nikisha nova
(Percentage)					
Carbohydrates	46	58	51	51	60-65
Protein	12	12	17	17	10-15
Fat	42	30	32	32	20-25

¹Obtained from Klicka et al. 1967, and U.S. Senate 1977.

Table 8

RELATIVE PERCENTAGE OF ESSENTIAL AMINO ACIDS
OF SOME PLANT PROTEINS COMPARED TO
AMINO ACIDS OF EGG PROTEIN¹

Amino Acids	Foodstuffs				
	Soybean meal, low fat	Whole Rice	Whole Wheat	Peanut Flour	Dried Roast Beans
Histidine ²	128	81	100	100	104
Threonine	80	78	67	57	79
Valine	76	88	62	66	78
Leucine	89	91	78	79	78
Isoleucine	97	84	64	66	89
Lysine	111	52	44	57	106
Methionine	53	106	78	25	62
Phenylalanine	95	89	91	88	89
Tryptophan	127	118	109	72	73

¹Calculated from FAO (1970a).

²Not generally considered essential for adults.

Table 9

ESTIMATED ESSENTIAL AMINO ACID REQUIREMENTS OF ADULTS^a

Essential Amino Acid	Men (mg/day)	Women (mg/day)
Histidine	700	450
Isoleucine	1100	620
Leucine	800	500
Methionine -		
in absence of cystine	1100	550
in presence of 810 mg cystine	200	(180)
Phenylalanine -		
in absence of tyrosine	1100	
in presence of 1100 mg tyrosine	300	(200)
Threonine	500	300
Tryptophan	250	160
Valine	800	650

^aThese estimates emphasize the upper range of individual requirements adapted from FAO (1973).

Lipids form an important part of the diet as a source of energy and for culinary reasons. Fats and oils are needed for cooking oils, spreads, dressings and flavorings. The lipid intake should not exceed 35% of the total energy supply. A daily consumption of approximately 60 grams is adequate for a person on a 2300 calorie diet (Wilks 1964).

Sufficient amounts of most vitamins and minerals can be supplied from plant sources. However, it is difficult to get sufficient riboflavin from plants, and vitamin B₁₂ does not occur in the plant kingdom. Among the minerals and trace elements, iodine may occur at suboptimal concentrations, and the low bioavailability of calcium, iron, zinc, and copper may render these deficient although they may nominally be present in sufficient amounts. Supplementation may be considered for these minor nutrients.

A 14-day cycle pure vegetarian menu was analyzed as a base to explore problem areas (Table 1). Dietary deficiencies for energy, protein, iron, vitamin B₁₂, and riboflavin are evident. A compilation of nutritional compositional data of ten different plant species is listed in Table 10. It further illustrates the difficulty of supplying sufficient energy intake while maintaining a balanced intake of nutrients. With implementation of modifications and further experimentation it can probably be developed into a satisfactory menu. The species diversity in this listing also recognizes the importance of some of the criteria mentioned earlier.

C. Oxygen Production and Carbon Dioxide Utilization

As actively-growing plants photosynthesize, carbon dioxide is fixed into organic carbon and oxygen is released. This oxygen can be utilized for human respiration if the atmosphere of the plant growing area is effectively integrated with the human habitation areas of the spacecraft. Plants can also help in maintenance of carbon dioxide levels in a regenerative life support

Table 10
Balanced Daily Diet Developed from Plant Sources¹

Plant Species	Quantity		General Composition				Minerals			Vitamins				
	Serving No.	Raw or Fresh Weight (g)	Cal.	Pro. (g)	Fat (g)	CHO (g)	Ca (mg)	Na (mg)	Fe (mg)	A (IU)	B ₁ (mg)	B ₂ (mg)	Nia (mg)	C (mg)
Soybean	4	360	468	39.5	20.8	38.8	245	5.4	9.1	27	.22	.09	.65	--
Split Pea	2	200	236	18.6	.8	48.0	99	26.0	3.7	80	.30	.08	1.80	--
Chick Pea	2	200(est)	226	18.0	4.0	48.0	88	25.0	4.0	32	.18	.10	1.24	--
Peanut	2	54	308	14.6	23.8	12.8	66	6.4	2.2	--	.12	.10	6.70	--
Rice	4	392	464	10.0	2.4	100.0	41	11.6	2.1	--	.36	.16	10.80	--
Wheat	4	272	456	16.8	2.0	96.0	50	4.1	4.3	--	1.32	.28	8.40	--
Corn	2	110	136	3.4	1.2	28.0	9	.4	.8	400	.24	.08	1.60	--
Sweet Potato	1	205	252	3.7	1.4	57.0	80	25.2	1.8	11,940	.14	.09	.09	26
Mustard Greens	1	70	15	1.6	.2	2.8	92	16.0	1.5	4,060	.06	.10	.40	34
Strawberry	1	149	55	1.2	.8	12.1	31	2.0	1.5	90	.04	.10	.90	88
TOTALS	23		2,615	127.5	57.4	443.5	801	122.1	31.0	16,629	2.98	1.18	33.39	148
% of RDA			97	288			~10	301-172		333	212	73	186	246

Total Volume: $\sim 2.7L \div 4 = 675$ ml per meal

Energy distribution: CHO: 67.8%, Protein: 19.5%; Fat 20.8%

¹Calculated from Adams (1975).

system. Carbon dioxide exhaled by humans in the spacecraft can be cycled to the plant growing area and assimilated into organic matter by the plants. However, it should be recognized that the CO_2 incorporation and the O_2 production by a growing crop will not necessarily represent the effective gas exchange in a regenerative life support system. Only exchange associated with digestible food obtained from each plant is useful. The oxygen that must be consumed during plant decomposition of this non-digestible fraction is equal in quantity to the oxygen released in the photosynthetic production of that fraction. Also, not until non-digestible plant parts have been decomposed (oxidized) to CO_2 and water will the life support system be truly regenerative. Therefore, plant species should preferably be selected to provide maximum edible and minimum non-edible biomass.

There is considerable variation among plants in their ability to produce oxygen but it is in direct proportion to the rate of net photosynthesis at any particular time. Thus, oxygen evolution can be assumed to be approximately proportioned to dry matter accumulation. The oxygen production in $\text{liters/m}^2/\text{day}$ is numerically equal to dry mass production in $\text{g/m}^2/\text{day}$ when the dry mass consists of sugars. This equity was utilized by Milov and Balakireva (1975) for estimating O_2 production from dry mass production.

It was found that 10 m^2 of plant growing area provided adequate oxygen for one person in a six months closed life support experiment (Gitel'son 1976). The minimum surface area required will vary according to the plant species being grown and the extent of surface area that can be continuously covered with photosynthesis tissue. Oxygen production and CO_2 incorporation by plants in the system could be balanced against human activity patterns in various way such as by:

- 1) Providing multiple plant growth compartment chambers so that a portion of the plant population would be photosynthesizing, while other portions are in darkness.
- 2) Altering radiation or temperature levels to moderate photosynthetic rates as needed.
- 3) Providing a system of gas storage.

Since maximum oxygen production and carbon dioxide utilization generally occurs during periods of maximum plant growth, frequent planting and harvesting will be required to assure desired gas exchange.

D. Water Recycling

Higher plants give off predictable quantities of water as transpiration, at rates that are considerably greater during light than during dark. The transpired water vapor can be used to maintain humidity in the plant growing areas or directed to other compartments of the spacecraft for humidity maintenance. Some of the water vapor may be condensed for use as potable water.

The rate of water release from plants can be controlled in several different ways to satisfy moisture and water demands within the spacecraft. Controls include varying the humidity level of the plant growing area, varying the intensity and length of the light period and amount of plant surface maintained.

It has been suggested (but not proven) that plants give off harmful substances, such as alkaloids, with the transpired water (Derendyayeva 1973) and thus water condensed from the air may need further purification before being used for drinking or in support of growth of other plants.

E. Waste Recycling

The possible use of higher plants for recycling human wastes within a regenerative life support system appears feasible (Berry et al. 1977; Furr et al. 1976);

Gordon 1978; Wallace et al. 1976). Lettuce has been successfully cultured on activated sludge after appropriate mineralization by microorganisms (Drake 1966). However, some problems have been encountered when Chinese cabbage, tampala and endive were cultured on activated sludge because certain nutritive elements were unavailable (Drake 1966). Russian studies (Tsvetkova et al. 1965) have shown that such problems can be overcome by proper mineral supplementation of the activated sludge.

F. Other Criteria

Plant species (and cultivars) should be selected with consideration for the following desirable, but not necessarily essential, morphological and physiological characteristics:

- 1) Short plants with high leaf density are desirable to maximize space utilization and light interception per unit volume.
- 2) Growth habit, determinate or indeterminate, for most efficient production over time.
- 3) Species and cultivars with a wide range of environmental tolerance should be selected for maximum flexibility in multiplant systems and tolerance to uncontrolled environmental extremes. This would include tolerance to a wide range of radiation, temperature, humidity, media moisture, nutrient, salt and pH levels.
- 4) Plants that accumulate or release toxic compounds into the atmosphere. On the other hand, species that effectively remove toxic compounds from the media or atmosphere should be utilized, providing the plant tissues do not accumulate any toxic components.
- 5) Species and cultivars should be selected that have maximum disease resistance in combination with high productivity.
- 6) Plants that produce irritating air borne pollen should not be utilized.
- 7) Self pollinating species are preferred to avoid problems of seedset.

III. LITERATURE CITED

- Adams, Catherine F. 1975. Nutritive value of American foods. ARS - USDA Agric. Handbook No. 456. Washington, D.C.
- Anderson, Michele, J. Rummel and S. Deutsch. 1979. Biospex: biological space experiments. A compendium of life sciences experiments carried on U.S. spacecraft. NASA Technical Memorandum 58217. Washington, D.C.
- Anonymous. 1965. Surveys of Soviet-bloc scientific and technical literature, Soviet bioastronautics and manned spaceflight, programs, organization, and personalities, comprehensive report, Aerospace Tech. Div. Rept., P-65-14. Library of Congress, Washington, D.C.
- Antipov, V.V., N.L. Delone, G.P. Parfyanov, V.G. Vystosky. 1964. Results of biological experiments carried out under conditions of flight ships Vostok with participation of cosmonauts A.G. Nikolayev, P.R. Popovich and V.F. Bykovsky. In: Highlights of Foreign Bioastronautics 1(13):5-15. Aerospace Med. Div. Brooks Air Force Base, TX.
- Berry, W.L., A. Wallace and O.R. Lunt. 1977. Recycling municipal wastewater for hydroponic culture. HortScience 12(3):186.
- Dadykin, V.P. 1968. Space plant-growing - USSR. Kosmiches koye Rasteniyevodstvo Znaniye Press. Biology series No. 1. Moscow. Translation Joint Pub. Res. Ser. N68-33260 Dept. Comm. Washington, D.C.
- Derendyayeva, T.A. 1977. Investigation of the possibility of using a condensate of transpiration moisture of Batata for cultivating plants in biological life support systems. Joint Pub. Res. Ser. L/6859 Dept. Comm. Washington, D.C. Moscow Kosmicheskaya Biologiya I Aviakosmicheskaya Meditsina 10(6):70-73 (1976).
- Dolan, Desmond D. 1973. Temperature photoperiod and light intensity effects on growth of Pisum sativum L. HortScience 8(4):330-331.
- Dole, S.H. 1964. The ecological complex in extraterrestrial bases. A paper presented to the third annual meeting of the Working Group on Extraterrestrial Resources on November 18-20, 1964. Rand Corp., Santa Monica, CA.
- Drake, C.L. 1966. Final technical report, study of life support systems for space missions exceeding one year in duration. 64-00504. Rpt. on NAS 2-3011 General Dynamics, Convair Div, San Diego, CA.
- Evans, L.T. ed. 1969. The induction of flowering, some case histories. Macmillan Co., Aust. Pty. Ltd., Victoria.
- Ewald, Ellen B. 1977. Recipes from a small planet. Ballantine Books, N.Y.
- Food and Agriculture Organization. 1967. Requirements of vitamin A, thiamin, riboflavine, and niacin. FAO Nutrition Meetings Rept. Series 41. Food and Agr. Organ., Rome.
- Food and Agriculture Organization. 1970a. Amino acid content of foods and biological data on proteins, FAO Nutritional Studies No. 24. Food and Agr. Organ., Rome.

- Food and Agriculture Organization. 1970b. Requirements of ascorbic acid, vitamin D, vitamin B-12, folate and iron. FAO Nutrition Meetings Rept. Series 47. Food and Agr. Organ. Rome
- Food and Agriculture Organization. 1973. Energy and protein requirements. FAO Nutrition Meetings Rept. Series 52, and WHO Tech. Rept. Series 522. Food and Agr. Organ. Rome.
- Furr, A.K., W.C. Kelly, C.A. Bache, W.H. Gutenmann and D.J. Lisk. 1976. Multi-element absorption by crops grown in pots on municipal sludge-amended soil. J. Agric. Food Chem. 24(4):889-892.
- Gaastra, P. 1959. Photosynthesis of crop plants as influenced by light, carbon dioxide, temperature and stomatal resistance. Meded. Landbouwhog. Wagen. Neth. 59(13):1-68.
- Gitel'son, I.I., B.G. Kovrov, G.M. Lisovskiy, Yu.N. Okladnikov, M.S. Rerberg, F.Ya. Sidko and I.A. Terskov. 1975. Problems of space biology, Vol. 28, Experimental Ecological Systems Including Man. Nauka Press, Moscow. Translation NASA Tech. Translation F-16993. Washington, D.C.
- Glembotskiy, Ya.L., A.A. Prokof'yeva-Belgovskaya, Z.B. Shamina, V.V. Khvostova, S.A. Valeva, N.S. Eyges and L.V. Nevzgodina. 1962. Influence of space-flight factors on heredity and development in Actinomycetes and higher order plants. In: N.M. Sisakyan, ed. Problems in Space Biology 1:259-271 USSR Acad. Sci. Pub. Howe Moscow. Translation NASA Tech. Translation F-174. Washington, D.C.
- Gordon, M.S. 1977. Biological recycling of dissolved nutrients in treated domestic wastewaters using hydroponic and aquacultural methods. 133-147. In: F.M. D'itri, ed. Wastewater Renovation and Reuse. Pollution Engineering and Technology Series Vol. III. Dekker Pub. Co., N.Y., NY.
- Hegsted, D.M. 1964. Proteins in space nutrition. 135-142. In: Conference on Nutrition in Space and Related Waste Problems. Univ. South Florida. Tampa, FL. Apr. 27-30. 1964. NASA SP-70 Sci. Tech. Inf. Div. Washington, D.C.
- Johnson, R.D. and C. Holbrow. 1977. Space settlements. A design study. SP-413 NASA Sci. Tech. Inf. Div. Washington, D.C.
- Klicka, M.V., H.A. Hollender and P.A. Lachance. 1967. Food for astronauts. J. Am. Dietet. Assoc. 51:238.
- Krauss, Robert W. 1962. Mass culture of algae for food and other organic compounds. Amer. J. Bot. 49(4):425-435.
- Kristoffersen, T. 1963. Interactions of photoperiod and temperature in growth and development of young tomato plants (Lycopersicum esculentum Mill.) Physiologia Plantarum Supp. I.
- Lebedeva, Ye. V. 1964. Characteristics of certain artificial substrates for use in a closed ecological system. In: N.M. Sisakyan and V.I. Yazdovskiy, ed. Problems in Space Biology 2:198-203. Translation NASA Tech. Translation 64-31578. Dept. Comm. Washington, D.C.

- Leung, W.T.W., R.R. Burum, F.H. Chang, M.N. Rao and W. Polacchi. 1972. Food composition table for use in East Asia. Dept. Health, Educ. and Welfare. Pub. 73-465. Washington, D.C.
- Mason, Robert M. and John C. Carden. 1979. Guiding the development of a controlled ecological life support system. Report on NASA/Ames Workshop Jan. 8-12, 1979. Grant NSG-2323. 55p. GA Inst. Tech., Atlanta, GA.
- Miller, R.L. and C.H. Ward. 1966. Algal bioregenerative systems. 185-222. In: Karl Kammermeyer, ed. Atmosphere in Space Cabins and Closed Environments. Meredith Pub. Co., N.Y., NY.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. 28-33. In: I.I. Gitel'son, ed. Problems of Creating Biotechnical Systems of Human Life Support. pp. 13-20. NASA Tech. Translation TT F-17533. Washington, D.C.
- National Academy of Sciences. 1980. Recommended Dietary Allowances, ninth edition. Nat. Acad. Sci.-Nat. Res. Coun. Publ. No. 2941. Washington, D.C.
- Ormrod, Douglas P. 1979. Personal Communications.
- Paul, A.A. and D.A. Southgate. 1978. The composition of foods. Her Majesty's Stationery Office, London, and Elsevier-North Holland Biomedical Press, N.Y.
- Rao, Rama Rao. 1965. Studies of the environmental factors controlling tipburn of lettuce. PhD thesis, University of Wisconsin, Madison, WI.
- Robertson, Laurel and Carol and Godfrey B. Slinders. 1978. Laurel's Kitchen, a handbook for vegetarian cookery and nutrition. Nilgiri Press, Berkeley, CA.
- Rusakova, G.G., V.V. Lebedeva, V.V. Smolyarova, M.V. Vil'yams, T.P. Alekhina and V.M. Simonov. 1975. The red beet for autotroph link in a biological life support system. In: I.I. Gitel'son, ed. Problems of Creating Biotechnical Systems of Human Life Support. Nauka Press, pp. 21-27. NASA Tech. Translation F-17533. Washington, D.C.
- Singh, O.S. 1978. Isolating the genetic potential of sugar beets seeds to germinate under osmotic and temperature stresses. Indian J. Ecol. 4(2):149-156.
- Space Science Board. 1969. Report of the Panel on Atmosphere Regeneration. National Academy of Sciences-National Research Council. Washington, D.C.
- Spurlock, Jack, et. al. 1979. Research planning criteria for regenerative life support systems applicable to space habitats. 13-30. In: John Billingham, ed. Space Resources and Space Settlements. SP-428. NASA Sci. Tech. Inf. Div. Washington, D.C.

- Taub, F.C. 1974. Closed ecological systems. *Ann. Rev. Ecol. Systemat.* 51: 139-167.
- Tsvetkova, I.V., Y.I. Shaydarov and V.M. Abramova. 1965. Special features of plant feeding under conditions of aeroponic cultivation for a closed system. *In: N.M. Sisakyan, ed. Problems of Space Biology*, 4:637-642. Translation NASA Tech. Translation F-368. Washington, D.C.
- United States Department of Agriculture. 1975. Composition of Foods Agric. Handbook 8. U.S. Govt. Print. Off. Washington, D.C.
- United States Department of Agriculture. 1979. Agricultural Statistics. U.S. Govt. Print. Off. Washington, D.C.
- United States Senate. 1977. Select committee on nutrition and human needs. Dietary goals for the United States. 2nd Ed., U.S. Gov. Print. Off., Washington, D.C.
- Wallace, A., P.M. Patel, W.L. Berry and O.R. Lunt. 1978. Reclaimed sewage water: a hydroponic growth medium for plants. *Resource Recovery and Conservation* 3:191-199.
- Ward, C.H., S.S. Wilks and H.L. Craft. 1963. Use of algae and other plants in the development of a life support system. *Am. Biol. Teacher* 25:512-521.
- Warrington, I.J., E.A. Edge and L.M. Green. 1978. Plant growth under high radiant energy fluxes. *Ann. Bot.* 42:1305-1313.
- Watt, Bernice K. and Annabel L. Merrill. 1963. Composition of Foods, USDA-ARS. Agric. Handbook No. 8. Washington, D.C.
- Wilks, S.S. 1959. Carbon monoxide in green plants. *Science* 129(3354):964-966.
- World Health Organization. 1962. Calcium requirements. WHO Tech. Rept. Series 230. World Health Organ. Geneva.

IV. RECOMMENDATIONS

A. Plant Species

Plant species for the regenerative life support systems were identified that would meet both the major nutritional needs of man and also represent uniquely different growth morphologies for system development. These species were designated as reference plants, representative of related species that might be utilized in an operational life support system.

Two groups of plants were selected for research study. The first group of eight plant species was designated for intensive study and are species that are commonly utilized food plants and could provide the major nutritional needs of man. Intensive research efforts should be undertaken to establish their baseline and optimum productivity in life support systems as detailed in the research priorities of Section C of these Recommendations.

The second group of six plant species was designated for exploratory study and were species for which there is less reported information or that have lower nutritive values but high psychological value. Research should be undertaken, and principally limited to, establishing their baseline productivity in a life support system and to developing culture procedures for possible use in a regenerative life support system.

It is intended that study of these two groups will provide the necessary information to develop an effective operational system for these or other food crops having similar or related growth requirements.

Reference Plants: Intensive Study

Wheat - Wheat was selected because of its high caloric density and because it is the basis for many different types of foods that can be produced with minimum processing. The edible proportion of the total biomass is high. Wheat

contains a high starch content, contains a reasonable amount of protein (to 14%), as well as phosphorus, iron, thiamin, niacin and fiber. It is recommended that dwarf cultivars be utilized. Use of wheat will be limited by the fact that photosynthesis ceases during seed maturation. Spring wheat cultivars should be selected in preference to winter wheat to avoid a requirement for vernalization.

Rice - Rice, as wheat, was selected for its high caloric density. It supplies slightly less protein (8%) but has a more nutritionally balanced protein than does wheat. Rice also supplies phosphorus, iron, thiamin and niacin. Research should be undertaken to determine if strains can be maintained in an indeterminate growth habit to increase productivity.

White Potatoes - White potatoes were selected because they are a high calorie food that can provide a wide variety of useful foodstuffs for man with minimum processing. The carbohydrate is not as dense as that of wheat and the protein concentration on a dry weight basis is similar to that of rice. White potatoes also provide a good source of vitamin C and potassium. In liquid or mist culture, tuber production will require unique mechanical systems for support. Researchers should explore the possibility of obtaining indeterminate growth habit of plants by early and frequent harvesting of the tubers to increase productivity.

Sweet Potato - Sweet potatoes have been selected because they are adapted to warm environments and also are a high calorie food. The food can be eaten with very little processing. The carbohydrate is about 30% more dense than that of white potatoes and the protein concentration is similar to that of white potatoes. Sweet potatoes provide potassium, vitamin A and vitamin C.

The leaves and young shoots of sweet potatoes are edible and may have limited use in the diet. Sweet potatoes also will require the development of unique mechanical support systems for the enlarged roots in liquid and mist culture systems. The vigorous vine-type growth of the stems may be a disadvantage for use of this species.

Soybeans - Soybeans were selected because they can serve as a major source of dietary protein. The seeds have greater concentrations of protein (45-50%) than any other common plant food. The essential amino acids are well balanced except that methonine tends to be low. The protein is highly digestible and has a wide range of utilization in the diet. There also is a significant amount of oil in soybeans that would be available for food preparation and other uses, but the oil must be processed before it can be used. Soybeans are a good source of phosphorus, iron, potassium and thiamin.

Soybeans have a reasonably well defined morphology that should permit efficient and easily manipulated growth in a space system. The inclusion of soybeans provides a plant type that is representative of other legumes that might be utilized in a life support system.

Peanuts - Peanuts also were selected as a major source of protein for the diet. The seeds contain about 25% protein. The essential amino acids are not as well balanced as in soybeans and methonine also tends to be low. The oil concentration is about 45%, or about twice that in soybeans. The oil can be expressed and utilized directly without processing. The seeds are another good source of phosphorus, iron, potassium, thiamin and niacin.

The growth habit of peanuts, involving seed production below the soil surface, may result in excessively complicated growing and harvesting procedures.

Lettuce - Lettuce was selected to provide a salad crop for variety in the diet. It provides significant quantities of vitamin A and C. It is a low growing crop with defined shape that has been cultivated extensively in controlled environments. Leaf, rather than heading types, are recommended to minimize production problems and permit multiple harvests from the same plants.

Sugar beets - Sugar beets were selected to provide sugar, both for the diet, and for possible culture of single-cell organisms as a secondary food source. Sugar

beets have a plant type that is representative of most other root crop species that might be utilized in a life support system as red beets, radishes, carrots, turnips and rutabagas. The beets can be eaten raw and the tops also are edible and may have use in the diet. The weight penalty for equipment to extract sugar from beets should be carefully determined.

Reference Plants: Exploratory Study

Taro - Taro was selected as a representative tropical root crop. Taro and other root crops (e.g. manioc) are utilized infrequently for food in temperate regions of the world. Investigations should be undertaken to determine if the yield of edible portion of this crop is comparable to that of white potatoes and sweet potatoes. The tops also might be utilized for food.

Winged beans - Winged beans were recommended for study because this species is under intensive study as a food source in many areas of the world. It is recognized as a good protein source, and both tops and roots can be eaten. It is adapted to warm temperatures, although additional study is needed to determine its edible productivity and acceptability as a food along with study of its growth under controlled environment conditions.

Broccoli - Broccoli was selected because it provides a rich source of vitamins A, B₁ (thiamin), B₂ (riboflavin), B₇ (niacin) and C. Broccoli has a growth habit similar to other Cruciferae (cabbage, kohlrabi, cauliflower, Chinese cabbage and Brussels sprouts), that might be utilized in a life support system. The green broccoli is recommended for study.

Strawberries - Strawberries were selected to include a fruit crop that would provide additional variety to the diet. The psychological value of this pleasing food may be very great in a life support system. Strawberries would provide large amounts of vitamins B₂, B₇ and C to the diet. As strawberries are low growing and can be maintained as single plants without runners, they could be

grown efficiently and manipulated easily in a space system. Day neutral varieties with continuous production would be preferable. There is a need to optimize production for strawberries in hydroponic culture.

Onions - Onions were selected to provide a commonly used source of seasoning for many foods. They provide no significant nutritional value to the diet. The growth habit of this plant will require different systems for support because there is neither a tap root (only a fibrous root), nor significant stem formation (only a bulb developing from fleshy leaf blades). Day neutral cultivars should be selected to insure bulb enlargement under long days.

Peas - Peas are included because dried peas are a rich protein source similar to that of peanuts and provide high amounts of methionine, which is low in soybeans and peanuts. Peas contain little fat, but contain relatively large quantities of minerals such as potassium, copper, iron, sulfur and phosphorus. Peas do not have a strong stem, so culture systems will have to be developed for effective shoot support for these plants. It may be possible to use cultivars having edible pods, but their harvestable seeds have lower nutritional value and lower productivity than conventional dried pea cultivars.

B. Growing Procedures

The following recommendations are intended as guidelines to indicate the environmental and cultural considerations that should be controlled and studied in the plant research. It is anticipated that these guidelines will be periodically updated as requirements for individual species are elucidated and new technology becomes available. The development of comprehensive quality assurance programs to detail seed handling procedures, replication, environmental measurement and calibration, data collection and statistical analysis should be undertaken during the first years of this program so that research undertaken in different laboratories can be compared effectively.

Temperatures - No particular guideline temperatures are proposed at this stage in the study of higher plants for regenerative life support systems. Temperature optima at different stages of development for each species must be established. Day-night temperature differentials should be studied to determine their optima under different radiation and humidity levels. Leaf temperatures should be measured so as to provide an accurate basis for air temperature control, especially when establishing effective radiation levels. Also, media or root temperatures should be monitored and possibly controlled separately from air temperatures to optimize growth of the plants.

Radiation - The radiation guideline is at least $325 \mu\text{E m}^{-2}\text{sec}^{-1}$ photosynthetic photon flux density (PPFD) in order that plant morphology and rates of growth at least approximate those of field growth of plants. The effects of radiation levels in excess of this level also should be explored for each species.

A $325\text{--}450 \mu\text{E m}^{-2}\text{sec}^{-1}$ level of radiation can be provided with 1500 ma fluorescent lamp (typically cool-white) supplemented with incandescent lamps in most standard reach-in and walk-in chambers. Levels to $600 \mu\text{E m}^{-2}\text{sec}^{-1}$ can be obtained with well constructed walk-in chambers. Higher levels of radiation will require the use of high intensity discharge (HID) lamps. Particular lamp types for high radiation levels need additional study, but phosphor coated metal-halide lamps alone or in combination with sodium lamps appear best for general high radiation level applications.

The use of long photoperiods of 18, 20 or 24 hour would permit irradiation with lower levels than if shorter photoperiods were utilized. The radiation provided to plants should be expressed as quantum-per-day to simplify comparison of radiation levels in studies.

Carbon dioxide - There is a need for continuous monitoring of carbon dioxide levels during growth rate investigations with plants because both the large increases in carbon dioxide when humans are present in the growth area and depletion when

there is a large photosynthetic surface, may greatly affect photosynthetic rates. Carbon dioxide control should be provided in chambers, where possible, with flexibility to control levels between 350 and 2000 ppm.

Atmospheric moisture - The need for humidity control in growth rate investigations with each plant species also is recognized. A relative humidity level of 70% is encouraged for general use at temperatures between 15 and 30°C during both the light and dark periods. It is recognized that this set RH level will provide different vapor pressure deficits at different temperatures, but this specific humidity level is encouraged until the requirements for particular species can be established more precisely.

Media - It was agreed that plant investigations involving productivity and nutrient interactions should be undertaken using hydroponic culture so that accurate knowledge of nutrient concentrations is obtained and the studies can be effectively duplicated. The hydroponic culture could involve use of liquid culture, liquid film technique, mist culture, or use of inert gravel or sand as supporting media.

Investigations involving aspects other than productivity and nutrient interactions, may be undertaken in peat-vermiculite or other solid media to simplify growing procedures providing growth rates and plant type in solid media are similar to those obtained in hydroponic culture.

pH and Conductivity - Solution pH must be audited regularly, and the pH adjusted at least once daily, or more frequently if containers and solution reservoirs are of small volume. The use of automatic pH controllers is recommended. The pH of the solution should be maintained within the range of 4-6.5 pH units. The conductivity of the solution should be monitored in order that a level is maintained that is between 1/2 and 2x the initial solution conductivity.

Seed Germination - Germination procedures must be developed for individual crop species to provide seedlings with enough elongation of the main axis to be effectively supported in hydroponic culture.

Plant Spacing - Optimum plant spacing must be determined from seeding to final harvest for each species. The spacing should be gradually increased with increasing plant size to assure maximum utilization of space. Optimum spacing will vary markedly among species as well as among cultivars.

Interval Between Plantings - The optimum interval between plantings to maintain a continuous food and oxygen supply will be determined separately for each species. It will be controlled by the length of the harvesting period, the length of storage period for the harvested crops, and the amount of photosynthesizing crop canopy needed at particular times to supply human O₂ requirements.

Length of Production Cycle - The length of the production cycle will vary for each crop species and for particular cultivars within a species.

Insect and Disease Control - A minimum amount of pesticide applications are recommended for insect and disease control. This is desirable to reduce the possibility of plant growth reductions as well as human toxicity from the applied pesticides. Construction of multiple plant compartments will minimize risk from insects and diseases. Plant growth compartments should be programmed to allow for short empty periods at intervals to aid in eradication of any possible insects and diseases before they can increase to serious levels. It is recommended that procedures be instituted to minimize the incidence of insects or diseases on seeds and other supplies at the initiation of each study. This will greatly reduce the risk of serious infestations.

Seed Maintenance and Storage - A supply of seed of each selected cultivar should be obtained from a primary producer so that the lot is known and additional seed can be obtained in the future from the same foundation stock. Seed should be

obtained and maintained at a location from which all scientists undertaking studies with a given species can utilize seed from the same stock. Seed should be sized to eliminate large and small seeds, and thus have a seed lot with maximum uniformity. Seed also should be carefully stored under the proper conditions of humidity and temperature so as to maintain maximum viability. The seed lot should be monitored for germination and vigor periodically. Small samples should be distributed in moisture proof containers for use in different laboratories with instructions on handling and storage.

C. Research Priorities

Research priorities have been divided into primary needs, secondary needs, and development needs. Primary needs are those factors required to provide a basis for utilizing, and determining the trade offs for use of higher plants in a regenerative life support system. Primary needs provide a basis for the functioning of other programs in the regenerative life support project; human requirements, waste management, and systems development. Secondary needs are those that are developed either from information generated from primary needs research or those required when the life support system is being integrated, i.e., growth of mixed species, recirculation of nutrients, utilization of waste from reference plants etc. Funding of secondary needs should be delayed 3-5 years after initiating funding of primary needs. Development needs involve primarily research that would increase the reliability of higher plant use in a regenerative life support system. These needs should be funded when there is agency commitment for the construction of regenerative life support systems for space use. Limited research might be undertaken in these areas earlier but should not preempt research efforts directed toward the primary and secondary needs that are outlined.

Primary Needs:

- 1) Develop a quality assurance program for the plant research. This should include:
 - a) seed maintenance and distribution
 - b) experimental design and replication
 - c) environmental measurement and control and instrument calibration
 - d) plant growth data collection and statistical analysis.
- 2) Establish and optimize the efficiency of production of digestible food, oxygen production and water recycling by reference plants on the basis of: volume and area of space required per unit time; labor requirements; weight of the plant growing system; electrical energy utilized; and purchase and maintenance costs of plant growing system. The research efforts should involve:
 - a) photosynthetically active radiation (PPFD) level, spectral balance and duration
 - b) light and dark temperature levels, shoot and root temperature differentials, temperature optima at different stages of growth
 - c) humidity levels during light and darkness
 - d) carbon dioxide concentration
 - e) nutrient balance and concentrations at different rates of growth and stages of maturity
 - f) watering procedures, amount and frequency
 - g) pH level and control
 - h) plant spacing at different stages of growth
 - i) selection of hydroponic culture method
 - j) germination procedures
 - k) plant support procedures

- 1) nutritional value of edible plant parts
 - m) cultivar selection
 - 3) Establish nutrient and volatile gas toxicity for individual species in a recirculating system. This will include:
 - a) heavy metal and volatile emissions from structural materials
 - b) volatile plant emissions and root exudations
 - 4) Determine utilization and tolerance of human and plant wastes by plants. This will include:
 - a) direct and partially-decomposed waste
 - b) nutrient composition and availability in wastes
 - c) pH control in wastes
 - d) elemental and organic toxicants in waste
- This research effort will require integration with microbiologists and engineers in the waste management program.

Secondary Needs:

- 1) Determine possible mutual toxicities between reference species when grown in the same system. This will include:
 - a) volatile emissions
 - b) exudations into nutrient media
 - c) toxicities in decomposed waste
- 2) Collation of data accumulation from reference plants to provide a data base for systems integration.

Development Needs:

- 1) Determine range of environmental, including substrate, conditions for efficient food production, oxygen production and water regeneration for each reference species including radiation, temperature, humidity, carbon dioxide, nutrients and pH.

- 2) Screen available germplasm and develop improved cultivars for greater efficiency in regenerative life support systems.
- 3) Determine capacity of reference species for removal of atmospheric contaminants from the spacecraft atmospheres.
- 4) Establish whether reference species can be grown effectively in low gravity environments.
- 5) Determine effectiveness of in vitro culture procedures for propagation of reference species.
- 6) Investigate use of plant hormones and growth regulators to increase the efficiency of reference species in a regenerative life support system.
- 7) Automation of planting, spacing and harvesting.

V. BIBLIOGRAPHY AND PRODUCTIVITY OF THE SELECTED PLANT SPECIES

The literature has been reviewed to obtain published information on controlled environment growth of the selected species. The data obtained is quite incomplete for there is limited data from controlled environments for many species. Also, it has often been impossible to translate the published data into meaningful data for use in life support systems. For example, plants in controlled environment studies have been maintained commonly in individual containers and therefore, unless container spacing is reported, the data can not be translated to production per-unit-area. Accurate comparison of productivity among different studies is limited by the fact that some studies are initiated from seeding in controlled environments while other studies are begun with transplants.

Literature citations for research undertaken with each species are provided in the following pages. The citations have been obtained both from scientists working with each of these plant species and through literature searches using available computer data bases. Some citations cover field research results when the data seemed particularly appropriate for the application of plants to life support systems. Most of the listed citations have been reviewed and only conclusive and appropriate literature citations have been included. Some citations were included even though not reviewed when the title or abstract indicated that the research would likely have application for life support systems.

A summary table of the available production information is provided (Table 11). The large variation in productivity among studies is a result both of differences in environmental conditions and also because some production data was based on growth for the total period from seeding to harvest and other data for only the period from transplanting to harvest. It is

difficult to estimate maximum productivity values for any of these species from this presently available data.

Table 11

PRODUCTION IN CONTROLLED ENVIRONMENTS*

Plant Species	Reference	Location	Growing Period (Days)	Production		
				Edible Dry Weight*** (g m ⁻² day ⁻¹)	Oxygen g m ⁻² day ⁻¹	Water kg m ⁻² day ⁻¹
Broccoli (<i>Brassica oleracea</i> var. <i>botrytis</i>)	(See below)**	Field	60 (Transplant)	1.31 (10.9)		
Lettuce (<i>Lactuca sativa</i>)	Milov & Novikova 1975 Hammer et. al. 1978 Tibbitts & Kozlowski 1980 Fontes 1973 (See below)**	Growth Chamb Growth Chamb Growth Chamb Greenhouse Field	22 (Transplant) 28 (Seeded) 28 (Seeded) 38 (Transplant) 60 (Seeded)	13.6 6.0 8.6 8.3 2.06 (4.9)	33.1	1.1
Onions (<i>Allium cepa</i>)	(See below)**	Field	110 (Seeded)	3.12 (10.9)		
Peanuts (<i>Arachis hypogaea</i>)	Milov & Novikova 1975 (See below)**	Growth Chamb Field	90 (No info) 120 (Seeded)	8.9 2.63 (94.4)	18.0	3.0
Peas, Dry (<i>Pisum sativum</i>)	(See below)**	Field	100 (Seeded)	1.68 (88.3)		
Rice (<i>Oryza sativa</i>)	Int. Rice Inst. 1976 Ishizuka 1979 (See below)**	Greenhouse Field Field	110 (Transplant) 110 (Transplant) 110 (Transplant)	8.4 7.6 3.25 (88.0)		
Soybeans (<i>Glycine max</i>)	Milov & Novikova 1975 Patterson et. al. 1977 Sionit & Kramer 1977 Raper & Thomas 1978 (See below)**	Growth Chamb Growth Chamb Growth Chamb Growth Chamb Field	80 (Transplant) 115 (Seeded) 120 (Seeded) 148 (Seeded) 120 (Seeded)	1.7 4.8 2.2 6.8 1.34 (90.0)	5.6	3.5
Strawberries (<i>Fragaria x ananassa</i>)	(See below)**	Field	365 (Plants)	0.12 (10.1)		
Sugar beets (<i>Beta vulgaris</i>)	Milov & Novikova 1975 (See below)**	Growth Chamb Field	100 (No info) 140 (Seeded)	32.4 5.22 (13.0)	62.6	2.0
Sweet Potatoes (<i>Ipomoea batatas</i>)	Milov & Novikova 1975 Huett 1975 Nikishanova 1977 (See below)**	Growth Chamb Field Growth Chamb Field	90 (No info) 147 (Transplant) 150 (No info) 120 (Transplant)	13.0 20.3 30.0 2.80 (29.4)	34.6	2.7
Taro (<i>Colocasia esculenta</i>)	(See below)**	Field	270 (No info)	2.21 (27.0)		
Wheat (<i>Triticum sativum</i>)	Gitel'son 1975 Gifford 1977 (See below)**	Growth Chamb Greenhouse Field	75 (Seeded) 130 (Seeded) 100 (Seeded)	16.4 9.4 3.85 (87.0)		
White Potatoes (<i>Solanum tuberosum</i>)	Milov & Novikova 1975 McCown & Kass 1977 Mendoza & Haynes 1976 (See below)**	Growth Chamb Growth Chamb Growth Chamb Field	80 (Transplant) 135 (Tubers) 120 (Tubers) 120 (Tubers)	19.0 14.4 6.7 4.30 (20.2)	40.8	2.5
Winged Beans (<i>Psophocarpus tetragonolobus</i>)	Nangju 1979	Field		1.1		

*Production from field environments is shown for comparison and when no controlled environment production was available for that species.

**Field production values determined from maximum yields published in agricultural statistics (USDA, 1979) and dry weights of produce reported in food handbook (USDA, 1975).

***Percent of fresh weight in parenthesis.

Broccoli

REFERENCE CITED

[illegible]

Broccoli

- Anderson, W.C. 1976. Tissue culture propagation of broccoli plants for use in F-1 hybrid seed production. HortScience 11(3 sect 2):297.
- Anderson, W.C., G.W. Meagher and A.G. Nelson. 1977. Cost of propagating broccoli plants through tissue culture. HortScience 12(6):543-544.
- Cutcliffe, J.A. 1975. Effect of plant spacing on single-harvest yields of several broccoli cultivars. HortScience 10(4):417-419.
- Maurer, A.R. 1976. Response of broccoli to five soil water regimes. Can. J. of Plant Sci. 56(4):953-959.
- Palevitch, D. and E. Pressman. 1973. Apex removal and single harvest yield of side shoots of broccoli. HortScience 8(5):411-412.
- Thompson, R. and H. Taylor. 1976. Plant competition and its implications for cultural methods in calabrese. J. of Hortic. Sci. 51(1):147-157.
- Weibe, H.J. 1975. The morphological development of cauliflower and broccoli cultivars depending on temperature. Scientia Horticulturae 3(1):95-101.

Lactuca sativa

44

Lettuce

- Bensink, J. 1971. On morphogenesis of lettuce leaves in relation to light and temperature. Meded. Landbouw. 71-15. Wageningen, Neth.
- Bierhuizen, J.F., J.L. Ebbens and N.C.A. Koomen. 1973. Effects of temperature and radiation on lettuce growing. Neth. J. Agric. Sci. 21:110-116.
- Enoch, H., I. Rylski and Y. Samish. 1970. CO₂ enrichment to cucumber, lettuce and sweet pepper plants grown in low plastic tunnels in a subtropical climate. Israel J. Agr. Res. 20(2):63-69.
- Fontes, M.R. 1973. Controlled-environment horticulture in the Arabian Desert at Abu Dhabi. HortScience 8(1):13-16.
- Hammer, P.H., T.W. Tibbitts, R.W. Langhans and J.C. McFarlane. 1978. Base-line growth studies of Grand Rapids lettuce (*Lactuca sativa* L.) in controlled environments. J. Amer. Soc. Hort. Sci. 103(5):649-654.
- Langhans, R.W. 1978. A Growth Chamber Manual. Cornell Univ. Press, Ithaca, NY.
- Milov, M.A. and G.M. Novakova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In I.I. Gitel'son (ed.) Problems of Creating Biotechnical Systems of Human Life Support. Nauka Press. Moscow Trans. NASA Tech. Trans. F-175333.
- Prince, R.P. and J.W. Bartok, Jr. 1978. Plant spacing for controlled environment plant growth. Trans. ASAE 21(2):332-336.
- Read, M. 1972. Growth and tipburn of lettuce: Carbon dioxide enrichment at different light intensity and humidity levels and rate of incorporation of carbon-14 assimilates into the latex. Ph.D. Thesis. Univ. of Wis. 145 pp.
- Soffe, R.W., J.R. Lenton and G.F.J. Milford. 1977. Effects of photoperiod on some vegetable species. Ann. App. Biol. 85(3):411-415.
- Tibbitts, T.W. and G. Bottenberg. 1976. Growth of lettuce under controlled humidity levels. J. Am. Soc. Hort. Sci. 101(1):70-73.
- Wiebe, H.-J. and H.-P. Lorenz. 1977. Influence of changing temperature and light-dependent temperature control on the growth of lettuce. Gartenbauw. 42(1):42-45.

Onions

REFERENCE CITED

[illegible]

Onions

Riekels, J.W. 1977. Nitrogen-water relationships of onions grown on organic soil. J. Am. Soc. Hort. Sci. 102(2):139-142.

Rogers, I.S. 1977. The influence of plant spacing on the frequency distribution of bulb weight and marketable yield of onions. J. Hort. Sci. 53(3): 153-161.

Peanuts

REFERENCE CITED

[illegible]

Peanuts

- Cox, F.R. 1979. Effect of temperature treatment on peanut vegetative and fruit growth. *Peanut Sci.* 6:14-17.
- Diener, U.L. and N.D. Davis. 1969. Production of aflatoxin on peanuts-D under controlled environments. *J. Stored Prod. Res.* 5(3):251-258.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In: I.I. Gitel'son (ed.) *Problems of Creating Biotechnical Systems of Human Life Support*. Nauka Press. Moscow Trans. NASA Tech. Trans. F-175333.
- Pallas, J.E., Jr. and Y.B. Samish. 1974. Photosynthetic response of peanut. *Crop Sci.* 14(3):478-482.
- Tsuno, Y. 1975. The influence upon the photosynthesis in several crop plants. *Proc. Crop Sci. Soc. Jpn.* 44(1):44-53.
- Wood, I.M.W. 1968. The effect of temperature at early flowering on the growth and development of peanuts (*Arachis hypogaea*). *Aust. J. Agric. Res.* 19: 241-251.
- Wynne, J.C., D.A. Emery and R.J. Downs. 1973. Photoperiodic responses in peanuts. *Crop Sci.* 13:511-514.
- Wynne, J.C. and D.A. Emery. 1974. Response of intersubspecific peanut hybrids to photoperiod. *Crop Sci.* 14(6):878-880.

Pisum sativum

50

Peas

- Dolan, D.D. 1972. Temperature, photoperiod and light intensity effects on growth of Pisum sativum L. Crop Sci. 12(1):60-62.
- Dolan, D.D. 1973. Temperature, photoperiod and light intensity effects on growth of Pisum sativum L. Northeast Region Plant Introduction Station. Geneva, NY. HortScience 8(4):330-331.
- McLean, L.A., F.W. Sosulski and C.G. Youngs. 1974. Effects of nitrogen and moisture on yield and protein in field peas. Can. Plant Sci. 54(2):301-305.
- Miller, D.G., C.E. Manning and I.D. Teare. 1977. Effects of soil water levels on components of growth and yield in peas. J. Amer. Soc. Hort. Sci. 102(3): 349-351.
- Summerfield, R.J., P.A. Huxley and F.R. Minchin. 1977. Plant husbandry and management techniques for growing grain legumes under simulated tropical conditions in controlled environments. Exp. Agric. 13(1):81-92.

Literature Citations

Rice

Oryza sativa

REFERENCE CITED		LOCA- TION	EXPERIMENT VARIABLES																DATA OBTAINED									
			Growth Chambers Greenhouse Field	Radi- ation	Temp- era- ture				Media									Production					Nutri- tion			Toxins		
																		Tops	Roots	Edible	Oxygen (carbon dioxide uptake)	Water	Carbohydrates	Proteins	Other	Human	Plant	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
Cock & Yoshida	1973	X							X											X								
Hanada	1974	X		X		X												X										
Haneda	1972	X				X																						
Hosoi	1975	X			X	X							X							X								
Int. Rice Inst.	1976	X	X	X	X	X	X		X	X						X			X		X	X	X		X	X	X	
Int. Rice Inst.	1976-1979	X	X	X	X	X	X		X	X	X		X			X			X	X	X		X		X	X	X	
Ishizuka	1979		X																	X								
Johnson & Diaz	1973	X									X									X								
Lin & Chen	1977		X			X													X		X	X						
Ojha & Pande	1976	X																	X									
Sarkar & Sircar	1975	X			X	X													X									
Sato	1971		X	X						X										X								
Sato et. al.	1974		X	X		X				X										X								
Sato & Takahashi	1971		X	X	X	X														X								
Shanghai Inst.	1977		X			X													X		X							
Singh	1973		X		X														X									
Singh & Pandya	1972		X		X														X									
Tsuno	1975			X						X											X	X						
Yoshida	1973		X			X													X									
Yoshida	1972		X	X	X	X		X		X									X		X	X						
Yoshida & Hara	1977		X		X		X												X		X							

Rice

- Cock, J.H. and S. Yoshida. 1973. Changing sink and source relations in rice (*Oryza sativa* L.) using carbon dioxide enrichment in the field. Soil Sci. Plant Nutr. 19(3):229-234.
- Hanada, K. 1974. Studies on branching habits in crop plants. Part 8. Varietal differences of tillering ability of rice plants under controlled conditions varying in light intensity and temperature. Proc. Crop Sci. Soc. Jpn. 43(1): 88-98.
- Haneda, E. 1972. Relationship between environmental factors and temperature of the rice plant. Part 1. Temperature of the rice plant in the phytotron. J. Yamagata Agric. For. Soc. 29:1-7.
- Hosoi, N. 1975. Effects of day length temperature and nitrogen level on the heading of rice plants under controlled environments. Proc. Crop Sci. Soc. Jpn. 44(4):382-388.
- International Rice Research Institute. 1976. Climate and Rice. Los Banos, Phillipines.
- International Rice Research Institute. 1976-1979. Annual Reports for 1975, 1976, 1977 and 1978. Los Banos, Phillipines.
- Ishizuka, V. 1979. The rice yield competition in Japan. Potash Rev. 1979(6): 1-10. Inter. Pot. Inst. Beine, Switzerland
- Johnson, L. and A. Diaz. 1973. A continuous rice production system. Agric. Mech. Asia 4(1):109-112.
- Lin, A.C. and C.S. Chen. 1977. Effect of various temperature durations on tillering and photosynthetic activities of rice. J. Agric. Assoc. China, 98:55-60.
- Ojha, T.P. and H.K. Pande. 1976. Study of growth dynamics of rice under controlled environment. Final technical report. Ind. Inst. Tech., Khragpur, India. 97 pp.
- Sarkar, K.K. and S.M. Sircar. 1975. Control of growth and reproduction in cultivated and wild rice varieties by light quality and dark period. Ann. Bot. 39(164):1063-1070.
- Sato, K. 1971. The development of rice grains under controlled environment. Part 2. The effects of temperature combined with air humidity and light intensity during ripening on grain development. Tohoku J. Agric. Res. 22(2): 69-79.
- Sato, K., K. Inaba and K. Honjo. 1974. The development of rice grains under controlled environment. IV. Properties of protein of rice kernels ripened under different temperatures combined with different light intensity and air humidity. Tohoku J. Agric. Res. 25(3/4):97-103.

- Sato, K. and M. Takahashi. 1971. The development of rice-M grains under controlled environment. Part 1. The effects of temperature, its daily range and photo period during ripening on grain development. Tohoku J. Agric. Res. 22(2):57-68.
- Shanghai Inst. Plant Physiol. Phytotron. 1977. The influence of high temperature on flowering and fruiting of early rice and its control. Part 3. The sensitivity of flowering and fruiting of early rice to high temperature injury. Acta Bot. Sin. 19(2):126-131.
- Singh, P.M. 1973. Effect of different levels of light intensity on vegetative growth of rice under controlled environment. Riso 22(2):97-103.
- Singh, P.M. and A.C. Pandya. 1972. Effect of different photoperiods of uniform daily amounts of radiation on vegetative growth of rice under controlled environment. Riso 21(3):239-245.
- Tsuno, Y. 1975. The influence of transpiration upon the photosynthesis in several crop plants. Proc. Crop Sci. Soc. Jpn. 44(1):44-53.
- Yoshida, S. 1973. Effects of temperature on growth of the rice plant (Oryza sativa L.) in a controlled environment. Soil Sci. Plant Nutr. 19(4):299-310.
- Yoshida, S. 1972. Physiological aspects of grain yield. Ann. Rev. Plant Physiol. 23:437-464.
- Yoshida, S. and T. Hara. 1977. Effects of air temperature and light on grain filling of an indica and a japonica rice (Oryza sativa L.) under controlled environment conditions. Soil Sci. Plant Nutr. 23(1):93-107.

Literature Citations

Soybeans

Glycine max

REFERENCE CITED		LOCA- TION	EXPERIMENT VARIABLES															DATA OBTAINED													
			Growth Chambers Greenhouse Field	Radi- ation	Temp- era- ture	Level Photoperiod Spectrum	Air Soil Leaf	Carbon dioxide	Humidity	Spacing & Growth Proc.	Media					Tissue culture	Chemical hormones	Production					Nutri- tion			Toxins					
											Type	Nutrients	pH	Soil potential & water stress	Tops			Roots	Edible	Oxygen (carbon dioxide uptake)	Water	Carbohydrates	Proteins	Other	Human	Plant					
																											1	2	3	4	5
Hofstra& Hesketh	1975	X	X					X													X	X									
Kaplan & Koller	1977	X																	X		X										
Lu & Yen	1975	X		X	X																										
Milov & Novikova	1975	X																		X	X	X				X	X				
Patterson & Kramer	1975	X	X	X																	X										
Patterson, et. al.	1977	X		X								X							X		X										
Peet, et. al.	1977	X										X									X										
Raper & Thomas	1978	X		X	X																										
Sionit & Kramer	1977	X																	X		X										
Summerfield, et. al.	1978																				X										
Thomas & Raper	1976	X		X																	X										
Tibbitts & Kozlowski	1980	X																	X												
Tsuno	1975		X							X												X	X								
Warrington & Mitchell	1975	X		X	X														X							X	X				
Warrington, et. al.	1977	X				X													X												
Woodward	1976	X	X																X		X										
Woodward & Begg	1976	X								X											X						X	X			

Soybeans

- Hofstra, G. and J.D. Hesketh. 1975. The effects of temperature and CO₂ enrichment on photosynthesis in soybean. In: R. Marcelle, ed. Environmental and biological control of photosynthesis. W. Junk. The Hague, Netherlands.
- Kaplan, S.L. and H.R. Koller. 1977. Leaf area and CO₂-exchange rate as determinants of the rate of vegetative growth in soybean plants. Crop Sci. 17(1): 35-38.
- Lu, Y.-C. and H. Yen. 1975. Photoperiod and temperature responses of soybean varieties observed in a phytotron. SABRAO J. 7(2):171-182.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In: I.I. Gitel'son (ed.) Problems of Creating Biotechnical Systems of Human Life Support. Nauka Press. Moscow Translation. NASA Tech. Trans. F-175333.
- Patterson, D.T. and P.J. Kramer. 1975. The relationship between light intensity and photosynthesis rate in cotton and soybeans from growth chambers and out-of-doors. Assoc. Southeast Biol. Bull. 22(2):72.
- Patterson, D.T., M.M. Peet and J.A. Bunce. 1977. Effect of photoperiod and size at flowering on vegetative growth and seed yield of soybean. Agron. J. 69(4): 631-635.
- Peet, M.M., D.T. Patterson and J.A. Bunce. 1977. The effect of increasing soybean density in controlled environments on their production and variability. Assoc. Southeast Biol. Bull. 24(2):76.
- Raper, D.D. and J.F. Thomas. 1978. Photoperiodic alteration of dry matter partitioning and seed yield in soybeans. Crop Sci. 18:654-656.
- Sionit, N. and P.J. Kramer. 1977. Effect of water stress during different stages of growth. Agron. J. 69:274-278.
- Summerfield, R.J., F.R. Minchin and E.H. Roberts. 1978. Realisation of yield potential in soyabean (Glycine max (L.) Merr.) and cowpea (Vigna unguiculata (L.) Walp.). Monograph, British Crop Protection Council 21:125-134.
- Thomas, J.F. and C.D. Raper, Jr. 1976. Photoperiodic control of seed filling for soybeans. Crop Sci. 16(5):667-672.
- Tibbitts, T.W. and T.T. Kozlowski. 1980. Growth of crop plants under high irradiation in the Wisconsin Biotron. Mimeograph. Univ. Wis. Biotron, Madison, WI.
- Tsuno, Y. 1975. The influence of transpiration upon the photosynthesis in several crop plants. Proc. Crop Sci. Soc. Jpn. 44(1):44-53.
- Warrington, I.J. and K.J. Mitchell. 1975. The suitability of three high intensity lamp sources for plant growth and development. J. Agric. Engng. Res. 20:295-302.

- Warrington, I.J., M. Peet, D.J. Patterson, J. Bunce, R.M. Haslemore and H. Hellmers.
1977. Growth and physiological responses of soybean under various thermoperiods.
Aust. J. Plant Physiol. 4:371-380.
- Woodward, R.G. 1976. Photosynthesis and expansion of leaves of soybean grown in
two environments. Photosynthetica 10(3):274-279.
- Woodward, R.G. and J.E. Begg. 1976. The effect of atmospheric humidity on the
yield and quality of soya bean. Aust. J. Agr. Res. 27(4):501-503.

Literature Citations

Strawberries

Fragaria x ananassa

REFERENCE CITED		LOCA- TION	EXPERIMENT VARIABLES																DATA OBTAINED												
			Growth Chambers Greenhouse Field			Radi- ation	Temp- era- ture			Media						Production					Nutri- tion			Toxins							
																Oxygen (carbon dioxide uptake)					Carbohy- drates		Proteins	Other	Human	Plant					
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
Albregts & Howard	1979		X								X								X		X										
Barritt	1974		X								X									X											
Bjurman	1974		X		X		X								X						X										
Bolton	1974		X																		X										
Brooks & Sargent	1976		X																		X										
Fujio & Amano	1974		X	X										X							X										
Hensley	1973		X								X								X												
Legeida et. al.	1976	X															X														
MacLachlan	1975		X								X																				
Moore et. al.	1975		X								X																				
Morris et. al.	1978		X								X																				
Morris et. al.	1979		X								X										X										
Tafazoli & Vince-Prue	1978		X		X														X												
Tafazoli & Shaybany	1978		X											X							X										

Strawberries

- Albregts, E.E. and C.M. Howard. 1978 (1979). Evaluation of plant density on strawberry plant growth and fruiting response. Proc. FL State Hort. Soc. 91(0):298-299.
- Barritt, B.H. 1974. Single harvest yields of strawberries in relation to cultivar and time of harvest. J. Am. Soc. Hort. Sci. 99(1):6-8.
- Bjurman, B. 1974. Strawberry yields in Sweden as influenced by cultivar, plant age and climate. Swed. J. Ag. Res. 4:219-231.
- Bolton, A.T. 1974. Yield of strawberries. Can. J. Plant Sci. 54(2):271-275.
- Brooks, A.H. and M.J. Sargent. 1976. Winter glasshouse strawberry production a sequential cropping program. Sci. Hortic. 4(4):353-360.
- Fujio, H. and T. Amano. 1974. Nutritional and physiological studies on betterment of quality of vegetable crops Part 2 effects of nutrients, heavy concentration of fertilizers, soil moisture and restriction of light on the yield and the quality of strawberry. Bull. Veg. Orn. Crops Res. Stn. Ser. C. (Kurume) 39-80.
- Hensley, J.G. 1973. Early runner production of strawberries as influenced by blossom removal and source of plants. HortScience 8(3):250.
- Legeida, V.S., B.A. Levenko, N.P. Berezenko, V.V. Liferova and G.R. Shchibrya. 1976. Production and study of callus tissue during culturing of mazzard cherry and strawberry anthers. Cytol. Genet. (Eng. Trans. Tsitol. Genet.) 10(6):5-9.
- MacLachlan, J.B. 1975. An estimate of the variability of the yield of individual strawberry plants and its significance for a breeding program. J. Hort. Sci. 50(4):399-403.
- Moore, J.N., G.R. Brown and H.L. Bowden. 1975. Evaluation of strawberry clones for adaptability to once-over mechanical harvest. HortScience 10(4):407-408.
- Morris, J.R., A.A. Kattan, G.S. Nelson and D.L. Cawthon. 1978. Developing a mechanical system for production harvesting and handling of processing strawberries. HortScience 13(3):260.
- Morris, J.R., S.F. Spayd, D.L. Cawthon, A.A. Kattan and G.S. Nelson. 1979. Strawberry, Fragaria ananassa, clonal fruit yield and quality response to hand picking prior to once-over machine harvest. J. Am. Soc. Hortic. Sci. 104(6):864-867.
- Tafazoli, E. and Daphne Vince-Prue. 1978. A comparison of the effects of long day and exogenous growth regulators on growth and flowering in strawberries, Fragaria x ananassa. Duch. J. Hort. Sci. 53(4):255-259.
- Tafazoli, E. and B. Shaybany. 1978. Influence of nitrogen, deblossoming, and growth regulator treatments on growth, flowering, and runner production of the 'Gem' everbearing strawberry. J. Am. Soc. Hort. Sci. 103(3):372-374.

Beta vulgaris

60

Sugar Beets

- Ford, M.A. and G.N. Thorne. 1967. Effect of CO₂ concentration on growth of sugar-beet, barley, kale and maize. *Ann. Bot. N.S.* 31(124):629-644.
- Hall, A.E. and R.S. Loomis. 1972. Photosynthesis and respiration by healthy and beet yellows virus-infected sugar beets (Beta vulgaris L.). *Crop Sci.* 12(5):566-572.
- Milford, G.F.J. and J.R. Lenton. 1976. Effect of photoperiod on growth of sugar beet. *Ann. Bot.* 40(170):1309-1315.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In: I.I. Gitelzon, ed. *Problems of Creating Biotechnical Systems of Human Life Support*. Nauka Press. Moscow Trans. NASA Tech. Trans. F-17533.
- Ohki, K. and A. Ulrich. 1973. Sugarbeet growth and development under controlled climatic conditions with reference to night temperature. *Amer. Sugar Beet Tech.* 17(3):270-279.
- Singh, O.S. 1978. Isolating the genetic potential of sugar beets seeds to germinate under osmotic and temperature stresses. *Indian J. Ecol.* 4(2):149-156.
- Snyder, F.W. 1975. Leaf and root accretion by sugarbeet seedlings in relation to yield. *Amer. Sugar Beet Tech.* 18(3):204-213.
- Ulrich, A. 1952. The influence of temperature and light factors on the growth and development of sugar beets in controlled climate environments. *Agron. J.* 44(2):66-73.
- Ulrich, A. 1955. Influence of night temperature and nitrogen nutrition on the growth, sucrose accumulation and leaf minerals of sugar beet plants. *Plant Phys.* 30(3):250-257.
- Ulrich, A. 1958. Effect of Climate on Sugar Beets Grown Under Standardized Conditions. *J. Am. Soc. Sugar Beet. Tech.* 10(1):1-23.
- Ulrich, A. 1961. Variety climate interactions of sugar beet varieties in simulated climates. *J. Am. Soc. Sugar Beet Tech.* 11:376-387.
- Ulrich, A. and K. Ohki. 1956. Hydrogen ion effects of the early growth of sugar beet plants in culture solution. *J. Amer. Soc. of Sugar Beet Tech.* 9:265-274.

Ipomoea batatas

62

Sweet Potato

- Bell, D.E. and R.G. Fuller. 1965. Physiological evaluation of multicellular plants in bioregenerative systems as a means of providing mans oxygen and food requirements. Report to Brooks Air Force Base, Aerospace Med. Div. Contract No. AF 41(609)-2391, Task 793001.
- Enyi, B.A.C. 1976. Analysis of growth and tuber yield in sweet potato (Ipomoea batatas) cultivars. J. Agr. Sci. 88:421-430.
- Huett, D.O. 1976. Evaluation of yield, variability and quality of sweet potato cultivars in sub-tropical Australia. Exper. Agric. 12(1):9-16.
- Lowe, S.B. and L.A. Wilson. 1974a. Comparative analysis of tuber development in six sweet potato (Ipomoea batatas (L.) Lam) cultivars. 1. Tuber initiation, tuber growth and partition of assimilate. Ann. Bot. 38(155):307-318.
- Lowe, S.B. and L.A. Wilson. 1974b. Comparative analysis of tuber development in six sweet potato (Ipomoea batatas (L.) Lam) cultivars. 2. Interrelationships between tuber shape and yield. Ann. Bot. 38(155):319-326.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In: I.I. Gitel'son (ed.) Problems of Creating Biotechnical Systems of Human Life Support. pp. 13-20. NASA technical translation TT F-17533. Washington, D.C.
- Nikishanova, T.I. 1978. Plants for space plantations. Piroda 10:105-117 (1977). Translation NASA Technical Memorandum. Washington, D.C.
- Plucknett, D.L. (ed.). 1970. Tropical root and tuber crops tomorrow. Vol. 1. University of Hawaii, Honolulu, HI.
- Tsuno, Y. 1975. The influence of transpiration upon the photosynthesis in several crop plants. Proc. Crop Sci. Soc. Jpn. 44(1):44-53.

Colocasia esculenta

64

Taro

- Bourke, R.M. and C.H. Perry. 1976. Influence of sett size on growth and yield of taro Colocasia-esculenta. Papua New Guinea Agric. J. 27(4):115-120.
- El-Habbasha, K.M., A.G. Behairy and M. Abol-Magd. 1976. The relation between method of propagation plant density and growth, mineral uptake and yield of dasheen Colocasia-antiquorum. Sci. Hortic. 4(1):15-22.
- Oliviera, J.F.S. 1974. The nutritional value of some foods consumed on San Tome Island. Ecol. Food and Nutr. 3(3):237-242.
- Plucknett, D.L. (ed.). 1970. Tropical root and tuber crops tomorrow. Vol. 1. Univ. Hawaii, Honolulu, HI.
- Raghu, A., T. Eadie and G.C. Llewellyn. 1978. Evaluation of potential for aflatoxin occurrence on celery, cauliflower, lettuce and taro root inoculated with Aspergillus flavus and A. parasiticus. J. of the Assoc. of Off. Anal. Chem. 61(4):998-1001.
- Sato, T., M. Kawai and T. Fukuyama. 1978. Matter production of taro plant Colocasia-esculenta. 1. Changes with growth in photosynthetic rate of single leaf. Jpn. J. Crop Sci. 47(3):425-430.
- Shaw, D.E. 1975. Illustrated notes on flowering, flowers, seed and germination in taro (Colocasia esculenta). Res. Bull., Dept. of Stock and Fish, Papua New Guinea, No. 13, 39-59.
- Tsuno, Y. 1975. The influence of transpiration upon the photosynthesis in several crop plants. Proc. Crop Sci. Soc. Jpn. 44(1):44-53.

Literature Citations

Wheat

Triticum sativum

REFERENCE CITED	LOCA- TION	EXPERIMENT VARIABLES																DATA OBTAINED											
		Growth Chambers Greenhouse Field	Radi- ation Level	Temp- era- ture Spectrum	Air Soil Leaf	Carbon dioxide	Humidity	Spacing & Growth Proc.	Media										Production				Nutri- tion			Toxins			
									Type	Nutrients	pH	Soil potential & water stress	Tissue culture	Chemical hormones	Tops	Roots	Edible	Oxygen (carbon dioxide uptake)	Water	Carbohydrates	Proteins	Other	Human	Plant					
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	
Ahmad	1973			X	X									X	X			X		X									
Anaya & Stolzy	1972	X													X	X													
Apel	1974	X							X											X									
Apel	1976	X																		X									
Austin & Edrich	1974	X			X													X											
Bagga & Rawson	1977	X				X												X		X									
Bird et. al.	1977	X				X			X																				
Evans	1978	X	X	X																X									
Focke	1975	X	X	X	X															X									
Ford & Thorne	1975	X		X		X														X									
Friend et. al.	1961	X			X													X											
Gifford	1977	X	X						X											X	X								
Gitel'son et. al.	1975	X																		X									
Keys et. al.	1977	X				X			X																				
Kolderup	1975a	X	X			X				X						X													
Kolderup	1975b	X				X										X				X					X				
Krenzer & Moss	1975	X				X			X											X									
Lipsett	1964	X											X							X							X		
Luxmoore et. al.	1972	X	X											X				X											
Macdowall	1971	X	X					X										X	X		X								
Morgan	1977	X														X			X										
Neales & Nicholls	1978	X							X										X										
Partridge & Shaykewich	1972	X				X							X		X					X						X			
Passera & Albuzio	1978	X														X			X		X								
Pomeroy & Fowler	1973	X				X																							
Rawson & Bagga	1979	X				X												X		X									
Sofield et. al.	1977a	X	X		X															X									
Sofield et. al.	1977b	X											X		X					X							X		
Sojka et. al.	1975	X					X											X											
Spiertz	1974	X	X	X		X														X									
Spiertz	1978	X	X	X		X						X						X	X						X	X			
Warrington et. al.	1977	X				X												X	X										
Wunsche	1973	X	X	X		X												X	X										
Yang & Jong	1972	X				X			X							X							X						
Vil'yams & Rumyantseva	1978	X																X	X		X								

Wheat

- Ahmad, Babaey. 1973. Investigation on development, growth and yield as a contribution to the study of productivity type in spring wheat, with special reference to photoperiod and temperature. Thesis, Justus-Liebig-Universitat, Giessen, Ger. Fed. Rep.
- Anaya, M.G. and L.H. Stolzy. 1972. Wheat response to different soil water-aeration conditions. *Soil Sci. Soc. Amer. Proc.* 36:485-489.
- Apel, P. 1974. Grain growth in spring wheat at different CO₂-concentrations. *Biochem. Physiol. Pflanzen*, 166(5/6):475-480.
- Apel, P. 1976. Studies on the possibility of influencing the yield of single ears in spring wheat. Experiments in growth chambers. *Kulturpflanze*, 24: 143-157.
- Austin, R.B. and J.A. Edrich. 1974. A comparison of six sources of supplementary light for growing cereals in glasshouses during winter time. *J. Ag. Eng. Res.* 19:339-345.
- Bagga, A.K. and H.M. Rawson. 1977. Contrasting responses of morphologically similar wheat cultivars to temperatures appropriate to warm temperate climates with hot summers a study in controlled environment. *Aust. J. Plant Physiol.* 4(6).
- Bird, I.F., M.J. Cornelium and A.J. Keys. 1977. Effects of temperature on photosynthesis by maize and wheat. *J. Exp. Bot.* 28(104):519-524.
- Evans, L.T. 1978. The influence of irradiance before and after anthesis on grain yield and its components in microcrops of wheat grown in a constant day-length and temperature regime. *Field Crops Res.* 1(1):5-19.
- Focke, R. 1975. Comparison of different types of lamps in greenhouses and growth chambers and their effect upon yield in spring barley and wheat. *Archiv. Zuchtungs.* 5(3):133-147.
- Ford, M.A. and G.N. Thorne. 1975. Effects of variation in temperature and light intensity at different times on growth and yield of spring wheat. *Ann. App. Biol.* 80(3):283-299.
- Friend, D.J.C., V.A. Helson and J.E. Fisher. 1961. The influence of the ratio of incandescent to fluorescent light on the flowering response of Marquis wheat grown under controlled conditions. *Can. J. Plant Sci.* 41(2):418-427.
- Gifford, R.M. 1977. Growth pattern, carbon dioxide exchange and dry weight distribution in wheat growing under differing photosynthetic environments. *Aust. J. Plant Physiol.* 4:99-110.
- Gitel'son, I.I., B.G. Kovrov, G.M. Lisovskiy, Yu.N. Okladnikov, M.S. Rerberg, F.Ya. Sidko and I.A. Terskov. 1975. Problems of Space Biology, Vol. 28, Experimental Ecological Systems Including Man. Nauka Press. Moscow, Trans. NASA Tech. Trans. F-16993. Washington, D.C.

- Keys, A.J., E.V.S.B. Sampaio, M.J. Cornelius and I.F. Bird. 1977. Effect of temperature on photosynthesis and photorespiration of wheat leaves. *J. Expt. Bot.* 28(104):525-533.
- Kolderup, F. 1975a. Effect of soil moisture, air humidity, and temperature on seed setting and ear size in wheat. *Acta Agr. Scand.* 25(2):97-102.
- Kolderup, F. 1975b. Effects of soil moisture and temperature on yield and protein in production in wheat. *Meld Nor Landbruk.* 54(28):1-18.
- Krenzer, E.G., Jr. and D.N. Moss. 1975. Carbon dioxide enrichment effects upon yield and yield components in wheat. *Crop Sci.* 15(1):71-74.
- Lipsett, J. 1964. The phosphorus content and yield of grain of different wheat varieties in relation to phosphorus deficiency. *Aust. J. Agric. Res.* 15:1-8.
- Luxmoore, R.J., R.E. Sojka and L.H. Stolzy. 1972. Root porosity and growth responses of wheat to aeration and light intensity. *Soil Sci.* 113(5):355-357.
- Macdowall, F.D.H. 1971. Growth kinetics of Marquis wheat II. Carbon dioxide dependence. *Can J. Bot.* 50:883-889.
- Morgan, J.M. 1977. Changes in diffusive conductance and water potential of wheat plants before and after anthesis. *Aust. J. Plant Physiol.* 4:75-86.
- Neales, T.F. and A.O. Nicholls. 1978. Growth responses of young wheat plants to a range of ambient CO₂ levels. *Aust. J. Plant Physiol.* 5:45-59.
- Partridge, J.R.D. and C.F. Shaykewick. 1972. Effects of nitrogen, temperature, and moisture regime on the yield and protein content of Neepawa wheat. *Can. J. Soil Sci.* 52(2):179-185.
- Passera, C. and A. Albuzio. 1978. Effect of salinity on photosynthesis and photorespiration of two wheat species (*Triticum durum* cv. PEPE 2122 and *Triticum aestivum* cv. Marzotto). *Can. J. Bot.* 56(1):121-126.
- Pomeroy, M.K. and D.B. Fowler. 1973. Use of lethal dose temperature estimates as indices of frost tolerance for wheat cold acclimated under natural and controlled environments. *Can. J. Plant Sci.* 53(3):489-494.
- Rawson, H.M. and A.K. Bagga. 1979. Influence of temperature between floral initiation and flag leaf emergence on grain number in wheat. *Aust. J. Plant Physiol.* 6:391-400.
- Sofield, I., L.T. Evans, M.G. Cook and I.F. Wardlaw. 1977. Factors influencing the rate and duration of grain filling in wheat. *Aust. J. Plant Physiol.* 4(5):785-797.
- Sofield, I., I.F. Wardlaw, L.T. Evans and S.Y. Zee. 1977b. Nitrogen, phosphorus and water contents during grain development and maturation in wheat. *Aust. J. Plant Physiol.* 4(5):799-810.

- Sojka, R.E., L.H. Stolzy and M.R. Kaufmann. 1975. Wheat growth related to rhizosphere temperature and oxygen levels. *Agron. J.* 67(5):591-596.
- Spiertz, J.H.J. 1974. Grain growth and distribution of dry matter in the wheat plant as influenced by temperature, light energy and ear size. *Neth. J. Agric. Sci.* 22(3):207-220.
- Spiertz, J.H.J. 1978. Grain production and assimilate utilization of wheat in relation to cultivar characteristics, climatic factors and nitrogen supply. *Agric. Res. Rept. Verslagen Landbou. Onder. No. 881, Wag. Neth.*
- Warrington, I.J., R.L. Dunstone and L.M. Green. 1977. Temperature effects at three development stages on the yield of the wheat ear. *Aust. J. Agric. Res.* 28(1):11-27.
- Wunsche, U. 1973. Influence of growth retarding substances on cereals. 2. Effects of temperature and light intensity on the influence of CC on tillering of barley and spring wheat. *Zeit. Acker u Pflanzen.* 138(2):129-136.
- Yang, S.J. and E. de Jong. 1972. Effect of aerial environment and soil water potential on the transpiration and energy status of water in wheat plants. *Agron. J.* 64(5):574-578.
- Vil'yams, M.V. and V.B. Rumyantseva. 1978. A study of gas exchange in the light and effectiveness of carbon dioxide utilization in biomass formation in wheat planting. *Inst. Med.-Biol. Probl., Moscow, USSR. Fiziol Rast. (Mosc.)* 25(4):681-687.

Solanum tuberosum

70

White Potato

- Frier, Vivienne. 1977. The relationship between photosynthesis and tuber growth in Solanum tuberosum L. J. Expt. Bot. 28(105):999-1007.
- Hammes, P.S., P.C. Nel and E.A. Beyers. 1974. Photoperiodic induction in potatoes. Crop Prod. 3:13-18.
- Krug, H. and W. Wiese. 1972. Effect of soil moisture on development and growth of the potato plant (Solanum tuberosum L.). Pot. Res. 15(4):354-364.
- Ku, Sun-Ben, Gerald E. Edwards and Champ R. Tanner. 1977. Effects of light, CO₂ and temperature on photosynthesis, O₂ inhibition of photosynthesis, and transpiration in Solanum tuberosum. Plant Physiol. 59:868-872.
- Lynch, D.R. and R.G. Rowberry. 1977. Population density studies with Russet Burbank. 2. The effect of fertilization and plant density on growth, development and yield. Amer. Potato 54(2):57-71.
- McCown, B.H. and I. Kass. 1977. Effect of production temperature of seed potatoes on subsequent yielding potential. Amer. Potato 54(6):277-287.
- McCown, B.G. and C.B. Tanner. 1980. Diurnal growth of potato tubers. Mimeograph available from Soils Department, Univ. of Wisconsin, Madison, WI.
- Mendoza, H.A. and F.L. Haynes. 1976. Variability for photoperiodic reaction among diploid and tetraploid potato clones from three taxonomic groups. Amer. Potato 53(9):319-332.
- Milov, M.A. and G.M. Novikova. 1975. Gas exchange and transpiration of higher plants in cultivation under artificial conditions. In: I.I. Gitel'son (ed.) Problems of Creating Biotechnical Systems of Human Life Support. pp. 13-20. NASA technical translation TT F-17533. Washington, DC.
- Nelson, S.H. and K.E. Hwang. 1975. Water usage by potato plants at different stages of growth. Amer. Potato 52(11):331-339.
- Smith, O. 1977. Potatoes: Production, Storage, Processing. 2nd Ed. Avi Pub. Co. Inc., Westport, CT.

Literature Citations

Winged Beans

Psophocarpus tetragonolobus

[illegible]

Winged Beans

- Harding, J., F.W. Martin and R. Kleiman. 1978. Seed protein and oil yields of the winged bean, Psophocarpus tetragonolobus, in Puerto Rico. Trop. Agric. 55(4):307-314.
- Herath, H.M.W. and D.P. Ormrod. 1978. Effects of temperature and photoperiod on winged beans, Psophocarpus tetragonolobus L. DC. Ann. Bot. 43:729-736.
- Karikari, S.K. and S. Oteng. 1977. The effect of staking on the growth and yield of the winged bean Psophocarpus tetragonolobus. Acta Hortic. (The Hague) 53:159-163.
- Khan, T.N. and W. Erskine. 1978. The adaption of winged bean (Psophocarpus tetragonolobus (L.) DC.) in Papua New Guinea. Aust. J. Agric. 29(2):281-289.
- Korte, R. 1974. The role of legumes in alleviating protein deficiency in Papua New Guinea. Sci. New Guinea 2(1):6-14.
- Levy, J. (ed.). 1978. The Winged Bean Flyer. Dept. of Agronomy, Univ. IL, Urbana, IL.
- Martin, F.W. and H. Delpin. 1978. Vegetables for the hot, humid tropics. Part 1. The winged bean, Psophocarpus tetragonolobus. Agric. Res. Ser. New Orleans, USA.
- Nangju, D. and J.P. Baudoin. 1979. Performance of winged bean (Psophocarpus tetragonolobus (L.) DC.) in Nigeria. HortScience 54(2):129-136.
- National Academy of Sciences. 1975. The winged bean: a high protein crop for the tropics. Special Report No. 17. National Academy of Sciences-National Research Council, Washington, D.C.
- Rachie, K.O. 1978. Productivity potentials of edible legumes in the lowland tropics 71-96. In: Gerald A. Jung (ed.) Crop Tolerance to Suboptimal Land Conditions. Special Publication, No. 32. American Soc. of Agron., Madison, WI.
- Thompson, A.E. 1978. Variation in Winged beans - an unexploited tropical food crop. (Abs.) HortScience 13(3 sect. 2):366.

Literature Citations

Cultural Procedures

REFERENCE CITED

REFERENCE CITED		LOCA- TION	EXPERIMENT VARIABLES															DATA OBTAINED													
			Growth Chambers Greenhouse Field	Radi- ation Level	Temp- era- ture Spectrum	Air Soil Leaf	Carbon dioxide	Humidity	Spacing & Growth Proc.	Media						Tissue culture	Chemical hormones	Production					Nutri- tion			Toxins					
										Type	Nutrients	pH	Soil potential & water stress	Tops	Roots			Edible	Oxygen (carbon dioxide uptake)	Water	Carbohydrates	Proteins	Other	Human	Plant						
																										1	2	3	4	5	6
Asher	1978	X	X									X	X																		
Asher & Edwards	1978													X																	
Asher & Edwards	1978													X																	
Asher <u>et al.</u>	1965											X		X																	
Berry	1978													X						X											
Berry & Lunt	1979													X						X											
Berry & Wallace	1979													X																	
Berry <u>et al.</u>	1977											X		X				X													
Bleak <u>et al.</u>	1974		X	X																	X										
Cooper	1979													X																	
Dalrymple	1973		X																		X										
Deutch & Rasmussen	1974					X														X											
Dormling <u>et al.</u>	1975	X																													X
Edwards & Ascher	1974	X										X		X																	
Enoch <u>et al.</u>	1970		X						X											X											
Ford & Thorne	1974									X																					
Furr <u>et al.</u>	1976											X		X																	
Gitel'son <u>et al.</u>	1976	X										X									X					X	X				
Hasegawa	1977						X															X	X								
Hoffman	1973		X							X						X				X		X									
Huxley & Summerfield	1976				X		X													X		X									
Imai & Murata	1976								X											X											
Imai & Murata	1977								X											X											
Imai & Murata	1978								X											X											
Kretschmer	1977				X															X											
Krizek et al.	1974	X			X		X		X											X											

Literature Citations

Cultural Procedures continued

REFERENCE CITED		LOCA- TION	EXPERIMENT VARIABLES																DATA OBTAINED										
			Growth Chambers Greenhouse Field		Radi- ation	Temp- era- ture				Carbon dioxide	Humidity	Spacing & Growth Proc.	Media				Soil potential & water stress	Tissue culture	Chemical hormones	Production					Nutri- tion			Toxins	
																				Oxygen (carbon dioxide uptake)					Carbohydrates	Proteins	Other	Human	Plant
																				Tops	Roots	Edible	Water						
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	
O'Leary & Knecht	1974	X									X							X											
McCree	1972	X			X													X			X								
McKinney & Menser	1976	X																										X	
Monteith	1976			X			X		X	X									X										
Moss & Loomis	1952		X																X			X							
Parkinson <u>et al.</u>	1974								X										X										
Rajan <u>et al.</u>	1971	X			X	X													X										
Rice	1974																		X									X	
Rozov	1976	X									X								X		X								
Rozov & Silyutina	1977	X											X						X										
Terskov <u>et al.</u>	1974	X									X										X								
Tibbitts	1979									X									X			X	X						
Tsvetkova <u>et al.</u>	1971										X		X																
Tsvetkova <u>et al.</u>	1970	X										X									X								
Von Volkenburgh & Davies	1977	X	X																X				X						
Wallace <u>et al.</u>	1978										X	X							X										
Wallace <u>et al.</u>	1978										X	X	X																
Warrington	1972	X								X				X		X			X				X						
Warrington <u>et al.</u>	1978	X		X															X										
Warrington & Mitchell	1975	X			X														X										
Warrington <u>et al.</u>	1976	X		X	X	X													X										
Wittwer	1970	X							X										X		X								
Wittwer & Robb	1964	X							X										X		X								
Wilkes	1964	X									X								X		X								

Cultural Procedures

- Asher, C.J. 1978. Natural and synthetic culture media for spermatophytes. In: M. Rechigl Jr. (ed.) CRC Handbook Series in Nutrition and Food. Vol. III. CRC Press, Inc. Cleaveland, OH.
- Asher, C.J. and D.G. Edwards. 1978. Critical external concentrations for nutrient deficiency and excess. In: A.R. Ferguson, R.L. Bielecki, I.B. Ferguson (eds.) Proceedings of the 8th International Colloquium on Plant Analysis and Fertilizer Problems. DSIR Information Series No. 134. Wellington Goot Printer, Wellington, N.Z.
- Asher, C.J. and D.G. Edwards. 1978. Relevance of dilute solution culture studies to problems of low fertility tropical soils. In: C.S. Andrew and E.J. Kamprath (eds.) Mineral Nutrition. CSIRO, Australia.
- Asher, C.J., P.G. Ozonne and J.F. Loneragan. 1965. A method for controlling the ionic environment of plant roots. Soil Sci. 100:149-156.
- Berry, Wade L. 1978. Comparative toxicity of VO_3^- , CrO_4^{2-} , Mn^{++} , Co^{++} , Ni^{++} , Cu^{++} , Zn^{++} and Cd^{++} to lettuce seedlings. In: P.C. Adrian and I. Brisbin (eds.) Environmental Chemistry and Cycling Processes Symposia. ERDA Symposium Series Conf. -760429. Tech. Inf. Cent. U.S. Dept. of Energy, Washington, D.C.
- Berry, Wade L. and O.R. Lunt. 1979. Tolerance to trace elements in plants. In: Miloslav Rechigl, Jr. (ed.) Handbook of Nutrition and Food CRC Press Inc.
- Berry, W.L. and A. Wallace (eds.) 1979. Trace element stress in plants. Effects and methodology. UCLA 12-1218 Lab. Nuclear Med. Rad. Biol. UCLA, Los Angeles, CA.
- Berry, Wade L., A. Wallace and O.R. Lunt. 1977. Recycling municipal wastewater for hydroponic culture. HortScience 12(3):186.
- Bleak, A.T., W. Keller and A.C. Hull, Jr. 1974. Range plant yield and species relationships in natural and partially controlled environments. Range Manage. 27(5):396-399.
- Dalrymple, D.G. 1973. Controlled environment agriculture: a global review of greenhouse food production. For. Agric. Econ. Rep. 89. Econ. Res. Ser. USDA, Washington, DC.
- Deutch, B. and O. Rasmussen. 1974. Growth chamber illumination and photomorphogenetic efficacy. Part 1. Physiological action of IR radiation beyond 750 nanometers. Physiol. Plant 30(1):64-71.
- Dormling, I., A. Gustafsson and G. Ekman. 1975. Growth disorders and phenotype variability in phytotron-cultivated barley. Hered. 79:255-271.

- Edwards, D.G. and C.J.A. Asher. 1974. The significance of solution flow rate in flowing culture experiments. *Plant Soil* 41:161-175.
- Enoch, H., Irene Rylski and Y. Samish. 1970. CO₂ enrichment to cucumber, lettuce and sweet pepper plants grown in low plastic tunnels in a subtropical climate. *Israel J. Agric. Res.* 20(2):63-69.
- Ford, M.A. and G.N. Thorne. 1974. Effects of atmospheric humidity on plant growth. *Ann. Bot.* 38(155):441-452.
- Furr, A.K., W.C. Kelly, C.A. Bache, W.H. Gutenmann and D.J. Lisk. 1976. Multielement absorption by crops grown in pots on municipal sludge-amended soil. *J. Agric. Food Chem.* 24(4):889-892.
- Gitel'son, I.I., I.A. Terskov, B.G. Kovrov, F.Ya. Sidko, G.M. Lisovsky, Yu.N. Okladnikov, V.N. Belyanin, I.N. Trubachov and M.S. Rerberg. 1976. Life support system with autonomous control employing plant photosynthesis. *Acta Astronaut.* 3(9-10):633-650.
- Hasegawa, S. 1977. Agro-climatological studies on C3-plants and C4 plants. 3. Transpiration rates and leaf temperatures. *J. Agric. Meteor.* 33(3):129-136.
- Hoffman, G.J. 1973. Humidity effects on yield and water relations of nine crops. *Trans. ASAE* 16(1):164-167.
- Huxley, P.A. and R.J. Summerfield. 1976. Effects of daylength and day/night temperatures on growth and seed yield of cowpea cv. K 2809 grown in controlled environments. *Ann. App. Biol.* 83(2):259-271.
- Imai, K. and Y. Murata. 1976. Effect of carbon dioxide concentration on growth and dry matter production of crop plants. 1. Effects on leaf area, dry matter, tillering, dry matter distribution ratio and transpiration. *Proc. Crop Sci. Soc. Jap.* 45(4):598-606.
- Imai, K. and Y. Murata. 1977. Effect of carbon dioxide concentration on growth and dry matter production of crop plants. 2. Specific and varietal differences in the response of dry matter production. *Jap. J. Crop Sci.* 46(2):291-297.
- Imai, K. and Y. Murata. 1978. Effect of carbon dioxide concentration on growth and dry matter production of crop plants. 3. Relationship between CO₂ concentration and nitrogen nutrition in some C3- and C4-species. *Jap. J. Crop Sci.* 47(1):118-123.
- Kretchmer, P.J., J.L. Ozbun, S.L. Kaplan, D.R. Laing and D.H. Wallace. 1977. Red and far-red light effects on climbing in Phaseolus vulgaris L. *Crop Sci.* 17:797-799.
- Krizek, D.T., W.A. Bailey, H. Klueter and R.C. Liu. 1974. Maximizing growth of vegetable seedlings in controlled environments at elevated temperature, light and carbon dioxide. *Acta Hort.* 39:89-102.
- O'Leary, J.W. and G.N. Knecht. 1974. Raising the maximum permissible air velocity in controlled environment plant growth chambers. *Physiol. Plant.* 32(2):143-146.

- McCree, K.J. 1972. The action spectrum, absorptance and quantum yield of photosynthesis in crop plants. *Agric. Meteorol.* 9:191-216.
- McKinney, H.H. and H.A. Menser. 1976. Chlorotic leaf markings in Betzes barley grown in a controlled environment chamber. Fleck-spot, physiologic leaf spot, genetic disorders of cereal plants. *Plant Dis. Rep.* 60(12):1017-1019.
- Monteith, J.L. (ed.). 1976. *Vegetation and the Atmosphere*. Academic Press, London. GB. 440 pp.
- Moss, R.A. and W.E. Loomis. 1952. Absorption spectra of leaves I. The visible spectrum. *Plant Physiol.* 27(2):
- Parkinson, K.J., H.L. Penman and E.B. Tregunna. 1974. Growth of plants in different oxygen concentrations. *J. Exp. Bot.* 25(84):132-145.
- Rajan, A.K., Brenda Betteridge and G.E. Blackman. 1971. Interrelationships between the nature of the light source, ambient air temperature and the vegetative growth of different species within growth cabinets. *Ann. Bot.* 35:323-343.
- Rice, E.L. 1974. Impact of allelopathy on agriculture. pp. 184-236 In: E.L. Rice (ed.) *Allelopathy*. Acad. Press, New York, NY.
- Rozov, N.F. 1976. Biology and productivity of vegetable plants in the phytotron in 3 year seedless cultivation. *Izv Timiryazev S-kh Adak* 4:141-151.
- Rozov, N.F. and Yu.I. Silyutina. 1977. Some regularities of root nutrition of plants in a phytotron under various daily doses. *Izv Timiryazev S-KH AKAD* 2:128-135.
- Terskov, I.A., I.I. Gitel'son, B.G. Kovrov, G.M. Lisovskiy, Y.N. Okladnikov, F.Y. Sid'ko, V.N. Belyanin and M.P. Shilenko. 1974. Experiment with the incorporation of vegetable plants in two semiclosed life support systems. Translation N74-28576:53-58. *Nat. Aero. Space Admin.* Wash. DC
- Tibbitts, T.W. 1979. Humidity and plants. *BioScience* 29:358-363.
- Tsvetkova, I.V., B.G. Gusarov, V.P. Zamota, Y.V. Maksimova and L.A. Filatkina. 1971. Problem of mineralization of vegetative waste products of a biocomplex by the method of thermal combustion. pp. 306-311 In: V.N. Chèrnigovskiy *Problems of Space Biology*, Vol. 16. "Nauka" Press, Moscow. Trans. TT F-719. N73-19111. *Nat. Aero. Space Admin.*, Washington, D.C.
- Tsvetkova, I.V., V.P. Zamota and E.V. Maksimova. 1970. Plant cultivation using keramzit in closed ecological systems. Trans. N70-23179:15-18. *Nat. Aero. Space Admin.*, Washington, D.C.
- Van Volkenburgh, E. and W.J. Davies. 1977. Leaf anatomy and water relations of plants grown in controlled environments and in the field. *Crop Sci.* 17(3): 353-358.

- Wallace, A., P.M. Patel, W. Berry, O.R. Lunt and R.T. Mueller. 1977. Hydroponic cultivation of higher plants. p. 139-147. In: F.M. D'Itri, ed. Wastewater Renovation and Reuse. Pollution Eng. and Tech. Series Vol. III. Dekker Pub. Co., New York, NY.
- Wallace, A., P.M. Patel, W.L. Berry and O.R. Lunt. 1978. Reclaimed sewage water: A hydroponic growth medium for plants. Resource Recovery and Conservation 3:191-199.
- Warrington, G.E. 1972. Transpiration and dry matter response to atmospheric humidity, matric suction, and fertility. Diss. Abs. Inter. B. 32(7):3735.
- Warrington, I.J., E.A. Edge and L.M. Green. 1978. Plant growth under high radiant energy fluxes. Ann. Bot. 42(182):1305-1313.
- Warrington, I.J. and K.J. Mitchell. 1975. The suitability of three high intensity lamp sources for plant growth and development. J. Agric. Eng. Res. 20:295-302.
- Warrington, I.J., K.J. Mitchell and G. Halligan. 1976. Comparisons of plant growth under four different lamp combinations and various temperature and irradiance levels. Agric. Meteorol. 16:231-245.
- Wittwer, S.H. 1970. Aspects of CO₂ enrichment for crop production. Trans. ASAE 13(2):249-251, 256.
- Wittwer, S.H. and W.M. Robb. 1964. CO₂ enrichment of greenhouse atmosphere for food crop production. Econ. Bot. 18(1):34-56.
- Wilks, S.S. 1964. Plant systems in long term flight nourishment sources. pp. 305-310. In: Natl. Aero. Space Admin. Conference on Nutrition in Space and Related Waste Problems. NASA SP-70 Washington, D.C.

VI. WORKSHOP PARTICIPANTS

ALFORD, DONALD K.
Department of Biology
Metropolitan State College
1006 11th Street
Denver, CO 80204

BERRY, WADE
Laboratory of Nuclear Medicine
& Radiation Biology
University of California
Los Angeles, CA 90024

BJORKMAN, OLLE
Department of Plant Biology
Carnegie Institute
Stanford, CA 94305

BLOOM, ARNOLD J.
Institute of Arctic Biology
University of Alaska
Fairbanks, AK 99701

HAMMER, P. ALLEN
Horticulture Department
Purdue University
W. Lafayette, IN 47907

HELLMERS, HENRY
Department of Botany
Duke University
Durham, NC 27706

HODGSON, RICHARD H.
USDA Weed Physiology &
Growth Regulation Research
Frederick, MD 21701

HOFF, JOHAN E.
Horticulture Department
Purdue University
W. Lafayette, IN 47907

HOSHIZAKI, TAK
Jet Propulsion Laboratory
4800 Oak Grove Drive
Pasadena, CA 91103

HOWE, JEAN M.
Department of Foods & Nutrition
Purdue University
W. Lafayette, IN 47907

HUFFAKER, RAY C.
Plant Growth Laboratory
University of California
Davis, CA 95616

JAFFE, MORDECAI
Department of Botany
Ohio University
Athens, OH 45701

KNOTT, WILLIAM
John F. Kennedy Space Center
J.F.K.S.C., FL 32899

KRIZEK, DONALD T.
Plant Stress Laboratory
USDA
Beltsville, MD 20705

LANGHANS, ROBERT W.
Department of Floriculture &
Ornamental Horticulture
Cornell University
Ithaca, NY 14853

McFARLANE, J. CRAIG
EPA Box 15027
Las Vegas, NV 89114

MITCHELL, CARY
Horticulture Department
Purdue University
W. Lafayette, IN 47907

NORTON, ROBERT
NW Washington Research &
Extension Unit
Washington State University
Mount Vernon, WA 98273

ORMROD, DOUGLAS P.
Department of Horticultural Science
University of Guelph
Guelph, Ontario
CANADA

PRINCE, RALPH
Agricultural Engineering Department
University of Connecticut
Storrs, CT 06268

RAINS, WILLIAM
Plant Growth Laboratory
University of California
Davis, CA 95616

RAPER, C. DAVID
Soil Science Department
NC State University
Raleigh, NC 27650

RAWLINS, STEPHEN
US Salinity Laboratory
4500 Glenwood Drive
Riverside, CA 92501

SALISBURY, FRANK B.
Department of Plant Science
UMC 48
Utah State University
Logan, UT 84322

TIBBITTS, T.W.
Department of Horticulture
University of Wisconsin
Madison, WI 53706

WARD, CALVIN H.
Department of Environmental Science
& Engineering
Rice University
Houston, TX 77001

ZILL, L. PETER
Ames Research Center
Moffett Field, CA 94035

1. Report No. NASA CP-2231		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle CONTROLLED ECOLOGICAL LIFE SUPPORT SYSTEM - USE OF HIGHER PLANTS				5. Report Date May 1982	
				6. Performing Organization Code	
7. Author(s) T. W. Tibbitts and D. K. Alford, Editors				8. Performing Organization Report No.	
9. Performing Organization Name and Address Dept. of Horticulture Dept. of Biology Univ. of Wisconsin Metropolitan State Madison, WI 53706 Denver, CO 80204				10. Work Unit No. T5425	
				11. Contract or Grant No. NSG-2405	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 24056				13. Type of Report and Period Covered Conference Publication	
				14. Sponsoring Agency Code 199-60-62	
15. Supplementary Notes Robert D. MacElroy, Contract Monitor, Mail Stop 239-10, NASA Ames Research Center, Moffett Field, CA 94035 (415) 965-5573 FTS 448-5573. The 3rd in a series of CELSS reports.					
16. Abstract This report summarizes the results of two workshops concerning the use of higher plants in Controlled Ecological Life Support Systems (CELSS). Criteria for plant selection were identified from these categories: food production, nutrition, oxygen production and carbon dioxide utilization, water recycling, waste recycling, and other morphological and physiological considerations. The 27 participants recommended types of plant species suitable for use in CELSS, growing procedures, and research priorities. Also included are productivity values for selected plant species.					
17. Key Words (Suggested by Author(s)) CELSS Regenerative Life Support System Crop Plants Plant Physiology				18. Distribution Statement Unclassified - Unlimited Subject Category 54	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 84	
				22. Price* A05	

National Aeronautics and
Space Administration

Washington, D.C.
20546

Official Business

Penalty for Private Use, \$300

SPECIAL FOURTH CLASS MAIL
BOOK

Postage and Fees Paid
National Aeronautics and
Space Administration
NASA-451



NASA

POSTMASTER: If Undeliverable (Section 158
Postal Manual) Do Not Return
